arXiv:1704.00329v1 [math.CT] 2 Apr 2017

ENRICHED DUALITY IN DOUBLE CATEGORIES: $\mathcal{V}$-CATEGORIES AND $\mathcal{V}$-COCATEGORIES

CHRISTINA VASILAKOPOULOU

Abstract. In this work, we explore a double categorical framework for categories of enriched graphs, categories and the newly introduced notion of cocategories. A fundamental goal is the establishment of an enrichment of $\mathcal{V}$-categories in $\mathcal{V}$-cocategories, which generalizes the so-called Sweedler theory relatively to an enrichment of algebras in coalgebras. The language employed is that of $\mathcal{V}$-matrices, and an interplay between the double categorical and bicategorical perspective provides a high-level flexibility for demonstrating essential features of these dual structures. Furthermore, we investigate enriched fibration aspects of categories of monads and comonads in fibrant double categories a.k.a. proarrow equipments, which lead to natural applications for their materialization as categories and cocategories in the enriched matrices setting.

Contents

1. Introduction 1
2. Background 3
2.1. Bicategories 3
2.2. Monoids and comonoids in monoidal categories 8
2.3. Universal measuring comonoid 10
2.4. Fibrations 12
3. Double categories 17
3.1. Background on fibrant double categories 17
3.2. Monads and comonads in double categories 21
3.3. Locally closed monoidal double categories 28
4. Enriched matrices, categories and cocategories 31
4.1. $\mathcal{V}$-matrices 32
4.2. $\mathcal{V}$-graphs and $\mathcal{V}$-categories 39
4.3. $\mathcal{V}$-cocategories 46
4.4. Enrichment of $\mathcal{V}$-categories in $\mathcal{V}$-cocategories 55
References 60

1. Introduction

In [HLFV15], an enrichment of the category of monoids in comonoids is established in a braided monoidal closed category $\mathcal{V}$ which is moreover locally presentable, induced by a generalization of Sweedler's *universal measuring coalgebra* $P(A, B)$ for algebras A, B [Swe69]. This constitutes an abstract framework for the so-called *Sweedler theory* of algebras and coalgebras [AJ13] in differential graded vector spaces, leading to an efficient formalism for the

bar-cobar adjunction in a broader effort to conceptually clarify the Koszul duality for (co)algebras [LV07].

The present work generalizes this result to its many-object setting; introducing the notion of a $\mathcal{V}$-*enriched cocategory* which reduces to a comonoid in $\mathcal{V}$ when its set of objects is singleton, we establish an enrichment of the category of $\mathcal{V}$-categories in $\mathcal{V}$-cocategories. This enrichment can be realized under the same assumptions on $\mathcal{V}$, and shares all fundamental characteristics with Sweedler theory for (co)monoids. In particular, the braided monoidal closed $\mathcal{V}$-**Cocat** acts on the monoidal $\mathcal{V}$-**Cat** via a convolution-like functor which exhibits its adjoint, the *generalized Sweedler hom* $T\colon \mathcal{V}\text{-}\mathbf{Cat}^{\mathrm{op}} \times \mathcal{V}\text{-}\mathbf{Cat} \to \mathcal{V}\text{-}\mathbf{Cocat}$ as the enriched-hom functor. Moreover, the enrichment in question is tensored and cotensored, via the *generalized Sweedler product* $\triangleright\colon \mathcal{V}\text{-}\mathbf{Cocat} \times \mathcal{V}\text{-}\mathbf{Cat} \to \mathcal{V}\text{-}\mathbf{Cat}$ and the action respectively.

The framework that brings all these structures together is that of a double category of $\mathcal{V}$-matrices, $\mathcal{V}$-$\mathbb{M}\mathbf{at}$. Its horizontal bicategory $\mathcal{V}$-**Mat** is well-studied even in the more abstract case of matrices enriched in bicategories [BCSW83], and our approach follows its one-object case as well as [KL01] in viewing $\mathcal{V}$-categories as monads therein and establishing important categorical properties. It turns out that working on the double categorical context offers a much clearer perspective for the categories of interest and their interrelations; for example, $\mathcal{V}$-functors are precisely *monad homomorphisms* (vertical monad maps in [FGK11]) in $\mathcal{V}$-$\mathbb{M}\mathbf{at}$ whereas only a special case of monad maps in $\mathcal{V}$-**Mat**.

For that purpose, we present a detailed framework for monads and comonads in arbitrary double categories, explore their (op)fibrational structure in the fibrant case [Shu08] and push the enrichment objective as far as possible. This way, demanding calculations involving enriched graphs, categories and cocategories (some of them found in [Vas14, §7]) are deduced from natural properties of monads and comonads in monoidal fibrant double categories, which are furthermore *locally monoidal closed*. By introducing this concept which endows the vertical and horizontal categories with a monoidal closed structure, we obtain an action of the category of monads on comonads which under certain assumptions induces the desired enrichment.

Finally, we are also interested in combining such an enrichment with the natural (op)fibred structure of monads and comonads. As a result, we first recall some general fibred adjunction results as well as some basic *enriched fibration* machinery from [Vas14] and subsequently apply it to the double categorical setting for the monoidal (op)fibrations of (co)monads, leading to respective results for $\mathcal{V}$-categories and $\mathcal{V}$-cocategories in $\mathcal{V}$-$\mathbb{M}\mathbf{at}$. It is expected that the abstract picture shall also contribute to applications for other (co)monads in double categories of similar flavor, indicatively for colored operads and cooperads.

A next step to this development will be to consider categories of *modules* and *comodules* for monads and comonads in double categories and look for similar enrichments, this time for the fibration $\mathcal{V}$-**Mod** $\to$ $\mathcal{V}$-**Cat** over an appropriately defined $\mathcal{V}$-**Comod** $\to$ $\mathcal{V}$-**Cocat** in $\mathcal{V}$-$\mathbb{M}\mathbf{at}$. This would again provide a many-object generalization of the enrichment of **Mod** in **Comod**, i.e. global categories of (co)modules for (co)monoids, established in [HLFV17].

A brief outline of the paper is as follows. Section 2 assembles all the necessary background material in order to make this paper as self-contained as possible. This includes known facts about bicategories, (co)monoids in monoidal categories and local presentability aspects, the theory of actions inducing enrichments, the universal measuring comonoid and the theory of fibrations and enriched fibrations. In order to restrain the length of the paper, we provide appropriate references whenever definitions and constructions are only sketched. In Section 3, we explore the double categorical framework for monads and comonads, considering monoidal and fibrant double categories also with a locally monoidal closed structure, and we combine it with the enriched fibration theory. Finally, Section 4 applies all the previous results to the double category $\mathcal{V}$-$\mathbb{M}\mathbf{at}$ of $\mathcal{V}$-matrices; by establishing necessary properties of monads ($\mathcal{V}$-categories) and comonads ($\mathcal{V}$-cocategories) therein, an enrichment between them is exhibited also on the (op)fibration level.

Acknowledgements. I would like to thank Martin Hyland for the insightful ideas that led to this work, as well as Ignacio López Franco for helpful discussions that affected many parts of this development.

2. Background

In this section, we gather all the necessary material for what follows, with the purpose of making this work as self-sufficient as possible. This material includes some basic bicategorical notions, constructions in monoidal categories with focus on the categories of monoids and comonoids and local presentability aspects, as well as parts of the theory of actions of monoidal categories inducing enrichment relations. Finally, we recall some results from related work concerning an enrichment of monoids in comonoids, and the recently introduced enriched fibration theory aimed to fit in a double categorical context which serves as the common framework for our objects of interest.

2.1. **Bicategories.** The original definition of a bicategory and a lax functor between bicategories can be found in Bénabou's [Bén67]. Other references, including the definitions of transformations and modifications are [Str96, Bor94a]. 2-categorical notions appear as 'strictified' versions of the bicategorical ones. Due to coherence for bicategories, we are often able to use 2-categorical machinery like pasting and mates correspondence directly in the weaker context. Categories of (co)monads in bicategories are carefully recalled; regarding 2-category theory, indicative references are [KS74, Lac10], whereas [Str72] presents the formal theory of monads in 2-categories.

Definition 2.1. A *bicategory* $\mathcal{K}$ is specified by objects $A, B, \ldots$ called *0-cells*, and for each pair of objects a category $\mathcal{K}(A, B)$, whose objects are called *1-cells* and whose arrows are called *2-cells*; vertical composition of 2-cells is denoted



and the identity 2-cell is $1_f \colon f \Rightarrow f \colon A \to B$. Moreover, for each triple of objects there is the *horizontal composition* functor $\circ : \mathcal{K}(B,C) \times \mathcal{K}(A,B) \to \mathcal{K}(A,C)$ which maps a pair of 1-cells (g,f) to $g \circ f = gf$ and a pair of 2-cells



Finally, for each object we have the *identity 1-cell* 1-cell $1_A : A \to A$.

The associativity and identity constraints are expressed via the *associator* with components invertible 2-cells $\alpha_{h,g,f} : (h \circ g) \circ f \xrightarrow{\simeq} h \circ (g \circ f)$ and the *unitors* by $\lambda_f : 1_B \circ f \xrightarrow{\simeq} f$, $\rho_f : f \circ 1_A \xrightarrow{\simeq} f$. The above are subject to coherence conditions: for $A \overset{f}{\to} B \overset{g}{\to} C \overset{h}{\to} D \overset{k}{\to} E$, the following commute

$$\begin{array}{ccc}
((k \circ h) \circ g) \circ f & \xrightarrow{\alpha_{k,h,g} * 1_f} & (k \circ (h \circ g)) \circ f \\
\downarrow{\scriptstyle \alpha_{kh,g,f}} & & \downarrow{\scriptstyle \alpha_{k,hg,f}} \\
(k \circ h) \circ (g \circ f) & & k \circ ((h \circ g) \circ f) \\
& {\scriptstyle \alpha_{k,h,gf}} \searrow \quad \swarrow {\scriptstyle 1_k * \alpha_{h,g,f}} & \\
& k \circ (h \circ (g \circ f)) &
\end{array} \tag{1}$$

$$\begin{array}{ccc}
(g \circ 1_B) \circ f & \xrightarrow{\alpha_{g,1_B,f}} & g \circ (1_B \circ f) \\
& {\scriptstyle \rho_g * 1_f} \searrow \quad \swarrow {\scriptstyle 1_g * \lambda_f} & \\
& g \circ f &
\end{array}$$

By functoriality of the horizontal composition we have $1_g \circ 1_f = 1_{g \circ f}$ and $(\beta' \cdot \beta) * (\alpha' \cdot \alpha) = (\beta' * \alpha') \cdot (\beta * \alpha)$, the latter known as the *interchange law*.

Given a bicategory $\mathcal{K}$, we may reverse only the 1-cells and form the bicategory $\mathcal{K}^{\mathrm{op}}$, with $\mathcal{K}^{\mathrm{op}}(A,B) = \mathcal{K}(B,A)$. We may also reverse only the 2-cells and form the bicategory $\mathcal{K}^{\mathrm{co}}$ with $\mathcal{K}^{\mathrm{co}}(A,B) = \mathcal{K}(A,B)^{\mathrm{op}}$. Reversing both 1-cells and 2-cells yields a bicategory $(\mathcal{K}^{\mathrm{co}})^{\mathrm{op}} = (\mathcal{K}^{\mathrm{op}})^{\mathrm{co}}$.

There are numerous examples of well-known and studied bicategories. Indicatively,

- $\mathbf{Span}(\mathcal{C})$ for any category $\mathcal{C}$ with pullbacks has objects the ones in $\mathcal{C}$, 1-cells spans $A \leftarrow M \to B$ and 2-cells span morphisms;
- $\mathbf{Rel}(\mathcal{C})$ for any regular category $\mathcal{C}$ is defined as $\mathbf{Span}(\mathcal{C})$ but with 1-cells relations $R \rightarrowtail X \times Y$, and composition is given by first taking the pullback and then performing epi-mono factorization;
- **BMod** has objects rings, 1-cells ring bimodules and 2-cells bimodule maps;
- $\mathcal{V}$-**Prof** has objects $\mathcal{V}$-categories, 1-cells $\mathcal{V}$-profunctors ($\mathcal{V}$-bimodules) $F \colon \mathcal{B}^{\mathrm{op}} \times \mathcal{A} \to \mathcal{V}$ and 2-cells appropriate $\mathcal{V}$-natural transformations.

Definition 2.2. Given bicategories $\mathcal{K}$ and $\mathcal{L}$, a *lax functor* $\mathscr{F} : \mathcal{K} \to \mathcal{L}$ consists of a mapping on objects $A \mapsto \mathscr{F}A$, a functor $\mathscr{F}_{A,B} : \mathcal{K}(A,B) \to \mathcal{L}(\mathscr{F}A, \mathscr{F}B)$ for every $A, B \in \mathcal{K}$, a natural transformation with components

$\delta_{g,f} : (\mathscr{F}g) \circ (\mathscr{F}f) \to \mathscr{F}(g \circ f)$ for any composable 1-cells, and a natural transformation with components $\gamma_A : 1_{\mathscr{F}A} \to \mathscr{F}(1_A)$ for every $A \in \mathcal{K}$.

The natural transformations γ and δ have to satisfy the following coherence axioms: for 1-cells $A \xrightarrow{f} B \xrightarrow{g} C \xrightarrow{h} D$, the following diagrams commute:

$$\begin{array}{ccc} (\mathscr{F}h \circ \mathscr{F}g) \circ \mathscr{F}f & \xrightarrow{\delta_{h,g}*1} & \mathscr{F}(h \circ g) \circ \mathscr{F}f \\ \downarrow{\scriptstyle \alpha} & & \downarrow{\scriptstyle \delta_{hg,f}} \\ \mathscr{F}h \circ (\mathscr{F}g \circ \mathscr{F}f) & & \mathscr{F}((h \circ g) \circ f) \\ \downarrow{\scriptstyle 1*\delta_{g,f}} & & \downarrow{\scriptstyle \mathscr{F}\alpha} \\ \mathscr{F}h \circ \mathscr{F}(g \circ f) & \xrightarrow[\delta_{h,gf}]{} & \mathscr{F}(h \circ (g \circ f)) \end{array} \tag{2}$$

$$\begin{array}{ccc} 1_{\mathscr{F}B} \circ \mathscr{F}f & \xrightarrow{\gamma_B*1} & \mathscr{F}(1_B) \circ \mathscr{F}f \\ \downarrow{\scriptstyle \lambda} & & \downarrow{\scriptstyle \delta_{1_B,f}} \\ \mathscr{F}f & \xleftarrow[\mathscr{F}\lambda]{} & \mathscr{F}(1_B \circ f) \end{array} \qquad \begin{array}{ccc} \mathscr{F}f \circ 1_{\mathscr{F}A} & \xrightarrow{1*\gamma_A} & \mathscr{F}f \circ \mathscr{F}(1_A) \\ \downarrow{\scriptstyle \rho} & & \downarrow{\scriptstyle \delta_{f,1_A}} \\ \mathscr{F}f & \xleftarrow[\mathscr{F}\rho]{} & \mathscr{F}(f \circ 1_A) \end{array} \tag{3}$$

If γ and δ are natural isomorphisms (respectively identities), then $\mathscr{F}$ is called a *pseudofunctor* or *homomorphism* (respectively *strict functor*) of bicategories. Similarly, we can define a *colax functor* of bicategories by reversing the direction of γ and δ, sometimes also called *oplax*. We obtain categories $\mathbf{Bicat}_l$, $\mathbf{Bicat}_c$, $\mathbf{Bicat}_{ps}$, $\mathbf{Bicat}_s$ with objects bicategories and arrows lax, colax, pseudo and strict functors respectively.

A *monoidal bicategory* $\mathcal{K}$ is a bicategory equipped with a pseudofunctor $\otimes\colon \mathcal{K} \times \mathcal{K} \to \mathcal{K}$ which is coherently associative and has an identity object I. The explicit definition with all appropriate diagrams can be found in any of the standard references, e.g. [Car95, §2.1]. In our examples, we will choose to establish monoidal structure of bicategories via the more general structure of a monoidal double category, as explained in Section 3; arguably, that technique is easier to apply given certain assumptions.

Definition 2.3. If $\mathscr{F}, \mathscr{G} : \mathcal{K} \to \mathcal{L}$ are two lax functors, a *lax natural transformation* $\tau : \mathscr{F} \Rightarrow \mathscr{G}$ consists of morphisms $\tau_A : \mathscr{F}A \to \mathscr{G}A$ in $\mathcal{L}$, along with natural transformations

$$\begin{array}{ccc} \mathcal{K}(A,B) & \xrightarrow{\mathscr{F}_{A,B}} & \mathcal{L}(\mathscr{F}A, \mathscr{F}B) \\ \downarrow{\scriptstyle \mathscr{G}_{A,B}} & \overset{\tau}{\Swarrow} & \downarrow{\scriptstyle \mathcal{L}(1,\tau_B)} \\ \mathcal{L}(\mathscr{G}A, \mathscr{G}B) & \xrightarrow[\mathcal{L}(\tau_A,1)]{} & \mathcal{L}(\mathscr{F}A, \mathscr{G}B) \end{array} \tag{4}$$

with components 2-cells $\tau_f\colon \tau_B \circ \mathscr{F}f \Rightarrow \mathscr{G}f \circ \tau_A$. This data is subject to standard axioms expressing the compatibility of τ with composition and units, using δ and γ of the lax functors.

A lax natural transformation τ is a *pseudonatural* transformation (respectively *strict*) when all the components τ_f of (4) are isomorphisms (respectively

identities). Also, an *oplax* natural transformation is equipped with a natural transformation in the opposite direction of (4). Note that between either lax or oplax functors of bicategories, we can consider both lax and oplax natural transformations.

Along with *modifications* between transformations (see [Bor94a]) we can form four functor bicategories of all different combinations of (op)lax functors, (op)lax natural transformations and modifications, e.g. $\mathbf{Bicat}_{l,l}(\mathcal{K},\mathcal{L})$, which contain $\mathbf{Bicat}_{ps,l}(\mathcal{K},\mathcal{L})$, $\mathbf{Bicat}_{ps,opl}(\mathcal{K},\mathcal{L})$ and $\mathbf{Hom}(\mathcal{K},\mathcal{L})$ of pseudofunctors and lax/oplax/pseudo natural transformations as sub-bicategories. When $\mathcal{L}$ is a 2-category rather than a bicategory, all these functor bicategories are also 2-categories.

Now a (strict) *2-category* is a bicategory in which all constraints are identities, *i.e.* $\alpha,\rho,\lambda = 1$. In this case, the horizontal composition is strictly associative and unitary and the axioms (1) hold automatically. Consequently, the collection of 0-cells and 1-cells form a category on its own.

Examples.

(1) **Cat** of (small) categories, functors and natural transformations;
(2) **MonCat** of monoidal categories, (strong) monoidal functors and monoidal natural transformations;
(3) $\mathcal{V}$-**Cat** of $\mathcal{V}$-enriched categories, $\mathcal{V}$-functors and $\mathcal{V}$-natural transformations for a monoidal category $\mathcal{V}$;
(4) $\mathbf{Fib}(\mathbb{X})$ and $\mathbf{OpFib}(\mathbb{X})$ of fibrations and opfibrations over $\mathbb{X}$, (op)fibred functors and (op)fibred natural transformations (see Section 2.4);
(5) $\mathbf{Cat}(\mathbb{E})$ of categories internal to $\mathbb{E}$, for a finitely complete category. Instances of this are ordinary categories ($\mathbb{E} = \mathbf{Set}$), double categories ($\mathbb{E} = \mathbf{Cat}$) and crossed modules ($\mathbb{E} = \mathbf{Grp}$).

The coherence theorems for bicategories and their homomorphisms are of great importance, and have been fundamental for the development of higher category theory. In particular, it is asserted that certain diagrams involving the constraint isomorphisms of bicategories will always commute. Coherence allows us to replace any bicategory with an appropriate strict 2-category.

Theorem 2.4. *Every bicategory is biequivalent to a 2-category.*

The proof is based on a bicategorical generalization of the Yoneda Lemma (see [Str80]), which states that the embedding

$$\begin{aligned} \mathcal{K} &\longrightarrow \mathbf{Hom}(\mathcal{K}^{\mathrm{op}}, \mathbf{Cat}) \\ A &\longmapsto \mathcal{K}^{\mathrm{op}}(A,-) \end{aligned}$$

is locally an equivalence, hence any bicategory $\mathcal{K}$ is biequivalent to a full sub-2-category of $\mathbf{Hom}(\mathcal{K}^{\mathrm{op}}, \mathbf{Cat})$.

We now turn to notions of monads and comonads in bicategories.

Definition 2.5. A *monad* in a bicategory $\mathcal{K}$ consists of an object B together with an endomorphism $t : B \to B$ and 2-cells $\eta : 1_B \Rightarrow t$, $m : t \circ t \Rightarrow t$ called the *unit* and *multiplication* respectively, such that the following diagrams

commute:



Equivalently, a monad in a bicategory $\mathcal{K}$ is a lax functor $\mathscr{F} : \mathbf{1} \to \mathcal{K}$, where $\mathbf{1}$ is the terminal bicategory with a unique 0-cell $\star$. This amounts to an object $\mathscr{F}(\star) = B \in \mathcal{K}$ and a functor $\mathscr{F}_{\star,\star} : \mathbf{1}(\star,\star) \to \mathcal{K}(B,B)$ which picks up an endoarrow $t : B \to B$. The natural transformations δ and γ of the lax functor give the multiplication and the unit of t

$$m \equiv \delta_{1_\star,1_\star} : t \circ t \to t \quad \text{and} \quad \eta \equiv \gamma_\star : 1_B \to t$$

and the axioms for $\mathscr{F}$ give the monad axioms for (t, m, η).

Remark 2.6. If $\mathscr{G} : \mathcal{K} \to \mathcal{L}$ is a lax functor between bicategories, the composite

$$\mathbf{1} \xrightarrow{\mathscr{F}} \mathcal{K} \xrightarrow{\mathscr{G}} \mathcal{L}$$

is itself a lax functor from $\mathbf{1}$ to $\mathcal{L}$, hence defines a monad. In other words, if $t : B \to B$ is a monad in the bicategory $\mathcal{K}$, then $\mathscr{G}t : \mathscr{G}B \to \mathscr{G}B$ is a monad in the bicategory $\mathcal{L}$, *i.e.* lax functors preserve monads.

Definition 2.7. A *(lax) monad functor* between two monads $t : B \to B$ and $s : C \to C$ in a bicategory consists of an 1-cell $f : B \to C$ between the 0-cells of the monads together with a 2-cell



satisfying compatibility conditions with multiplications and units:





If the 2-cell ψ is in the opposite direction, and the diagrams are accordingly modified, we have a *colax* monad functor (or monad *opfunctor*) between two monads. Along with appropriate notions of monad natural transformations (see [Str72]), we obtain a bicategory $\mathbf{Mnd}(\mathcal{K}) \equiv [\mathbf{1}, \mathcal{K}]_l$ by the correspondence with lax functors from the terminal bicategory.

Dually to the above, we have the following definitions.

Definition 2.8. A *comonad* in a bicategory $\mathcal{K}$ consists of an object A together with an endoarrow $u : A \to A$ and 2-cells $\Delta : u \Rightarrow u \circ u$, $\varepsilon : u \Rightarrow 1_A$ called the *comultiplication* and *counit* respectively, such that the following commute



Notice that a comonad in the bicategory $\mathcal{K}$ is precisely a monad in the bicategory $\mathcal{K}^{\mathrm{co}}$; along with oplax comonad functors and comonad natural transformation, we have the bicategory $\mathbf{Cmd}(\mathcal{K}) = [\mathbf{1}, \mathcal{K}]_c$.

2.2. **Monoids and comonoids in monoidal categories.** Standard references for the theory of monoidal categories, as well as monoids and comonoids, are for example [JS93, Str07]. Local presentability properties [AR94] follow the development in [Por08b].

Suppose $(\mathcal{V}, \otimes, I)$ is a monoidal category. A *monoid* is an object A equipped with a multiplication and unit $m : A \otimes A \to A \leftarrow I : \eta$ that satisfy usual associativity and unit laws; along with monoid morphisms that preserve the structure in that $f \circ (m \otimes m) = m' \circ f$ and $f \circ \eta = \eta'$, they form a category $\mathbf{Mon}(\mathcal{V})$. Dually, we have *comonoids* $(C, \Delta\colon C \to C \otimes C, \epsilon\colon C \to I)$ whose category is denoted by $\mathbf{Comon}(\mathcal{V})$. Both these categories are monoidal only if $\mathcal{V}$ is braided, and they also inherit the braiding or symmetry from $\mathcal{V}$.

Remark 2.9. For any object B in a bicategory $\mathcal{K}$, the hom-category $\mathcal{K}(B, B)$ is equipped with a monoidal structure induced by the horizontal composition of the bicategory, namely $f \otimes g = g \circ f$ and $I = 1_B$. Then, a monoid in $(\mathcal{K}(B, B), \circ, 1_B)$ is precisely a monad in $\mathcal{K}$ and dually, a comonad $u : A \to A$ in a bicategory $\mathcal{K}$ is a comonoid in the monoidal $\mathcal{K}(A, A)$.

Lax monoidal functors between monoidal categories induce functors between their category of monoids, as the following well-known proposition demonstrates.

Proposition 2.10. *If $F\colon \mathcal{V} \to \mathcal{W}$ is a lax monoidal functor, with structure maps $\phi_{A,B}\colon FA \otimes FB \to F(A \otimes B)$ and $\phi_0\colon I \to F(I)$, it induces a map between their categories of monoids*

$$\mathbf{Mon}F\colon \mathbf{Mon}(\mathcal{V}) \to \mathbf{Mon}(\mathcal{W})$$

by $(A, m, \eta) \mapsto (FA, Fm \circ \phi_{A,A}, F\eta \circ \phi_0)$. Dually, colax functors induce maps between the categories of comonoids.

It is also well-known that due to doctrinal adjunction, colax monoidal structures on left adjoints correspond bijectively to lax monoidal structures on right adjoints between monoidal categories; this generalizes to parametrized adjunctions, e.g. [Vas14, 3.2.3] or for higher dimension in [DS97, Prop. 2].

When $\mathcal{V}$ is braided monoidal closed, the tensor product is strong monoidal via isomorphisms $A \otimes B \otimes A' \otimes B' \xrightarrow{\sim} A \otimes A' \otimes B \otimes B'$, $I \xrightarrow{\sim} I \otimes I$. Therefore the

internal hom functor $[-,-]\colon \mathcal{V}^{\mathrm{op}} \times \mathcal{V} \to \mathcal{V}$ obtains a lax monoidal structure as its parametrized adjoint. The induced functor between the monoids is denoted

$$\mathbf{Mon}[-,-]\colon \mathbf{Comon}(\mathcal{V})^{\mathrm{op}} \times \mathbf{Mon}(\mathcal{V}) \to \mathbf{Mon}(\mathcal{V}); \tag{5}$$

for C a comonoid and A a monoid, $[C, A]$ has the *convolution* monoid structure.

For any $\mathcal{V}$, there exist forgetful $S\colon \mathbf{Mon}(\mathcal{V}) \to \mathcal{V}$, $U\colon \mathbf{Comon}(\mathcal{V}) \to \mathcal{V}$; when these have a left or right adjoint respectively, these are called the *free monoid* and *cofree comonoid* functors. Evidently, the free monoid construction is quite frequent.

Proposition 2.11. *Suppose that $\mathcal{V}$ is a monoidal category with countable coproducts which are preserved by $\otimes$ on either side. The forgetful $\mathbf{Mon}(\mathcal{V}) \to \mathcal{V}$ has a left adjoint L, and the free monoid on an object X is given by*

$$LX = \coprod_{n\in\mathbb{N}} X^{\otimes n}.$$

On the other hand, the existence of the cofree comonoid is more problematic, and has been studied from various authors mainly in the context of vector spaces or modules over a commutative ring. We are interested in Porst's approach [Por08a, Por08b] which in particular focuses on local presentability properties inherited from $\mathcal{V}$.

Recall that an *accessible* category $\mathcal{C}$ is one with a small set of κ-presentable objects C (i.e. $\mathcal{C}(C,-)$ preserves κ-filtered colimits) such that every object in $\mathcal{C}$ is the κ-filtered colimit of presentable objects, for some regular cardinal κ. A functor between accessible functors is *accessible* if it preserves κ-filtered colimits. A *locally presentable* category is an accessible category which is cocomplete. More on the theory of locally presentable categories can be found in the standard [AR94]. An important fact is that any cocontinuous functor from a locally presentable category has a right adjoint; this can be seen as a corollary to the following adjoint functor theorem, since presentable objects form a small dense subcategory of $\mathcal{C}$.

Theorem 2.12. [Kel05, 5.33] *If the cocomplete $\mathcal{C}$ has a small dense subcategory, every cocontinuous $S : \mathcal{C} \to \mathcal{B}$ has a right adjoint.*

Proposition 2.13. [Por08b, 2.6-2.7] *Suppose $\mathcal{V}$ is a locally presentable monoidal category, such that $\otimes$ preserves filtered colimits in both variables.*

(1) $\mathbf{Mon}(\mathcal{V})$ *is finitary monadic over $\mathcal{V}$ and locally presentable.*
(2) $\mathbf{Comon}(\mathcal{V})$ *is a locally presentable category and comonadic over $\mathcal{V}$.*

The proof uses categories of *functor algebras* and *coalgebras* $\mathbf{Alg}F$ and $\mathbf{Coalg}F$ for any endofunctor $F\colon \mathcal{C} \to \mathcal{C}$, namely objects A equipped with plain arrows $FA \to A$ or $A \to FA$ in $\mathcal{C}$ and morphisms that commute with them. Their most important properties are the following.

Lemma 2.14. *For any endofunctor $F\colon \mathcal{C} \to \mathcal{C}$,*

(1) $\mathbf{Alg}F \to \mathcal{C}$ creates all limits and those colimits preserved by F;
(2) $\mathbf{Coalg}F \to \mathcal{C}$ creates all colimits and those limits preserved by F;
(3) if $\mathcal{C}$ is locally presentable and F preserves filtered colimits, $\mathbf{Alg}F$ and $\mathbf{Coalg}F$ are locally presentable.

Notably, these categories can be expressed as specific *inserters* $\mathbf{Alg}F = \mathbf{Ins}(F, \mathrm{id}_{\mathcal{C}})$ and $\mathbf{Coalg}F = \mathbf{Ins}(\mathrm{id}_{\mathcal{C}}, F)$ and Item 3 then follows from the more general 'Limit Theorem' [MP89, 5.1.6].

For the endofunctors with mappings $T_+(C) = (C \otimes C) + I$ and $T_\times(C) = (C \otimes C) \times I$, $\mathbf{Mon}(\mathcal{V})$ is a complete full subcategory of the locally presentable and finitary monadic over $\mathcal{V}$ category $\mathbf{Alg}T_+$, and $\mathbf{Comon}(\mathcal{V})$ is a cocomplete full subcategory of the locally presentable and comonadic over $\mathcal{V}$ $\mathbf{Coalg}T_\times$.

Specifically for comonoids, local presentability is deduced by expressing it as an equifier of a triple of natural transformations between accessible functors. Then comonadicity follows: in the commutative triangle

$$\begin{array}{ccc} \mathbf{Comon}(\mathcal{V}) & \hookrightarrow & \mathbf{Coalg}T_\times \\ & \overset{U}{\searrow} & \downarrow \\ & & \mathcal{V} \end{array} \tag{6}$$

where all categories are locally presentable, both forgetful functors to $\mathcal{V}$ have a right adjoint by Theorem 2.12, since they are cocontinuous. Moreover, the right leg is comonadic by Item 2, and the inclusion preserves and reflects all limits. Therefore it creates equalizers of split pairs and so does U, which then satisfies the conditions of Precise Monadicity Theorem. In particular, the existence of the *cofree comonoid functor* $R : \mathcal{V} \to \mathbf{Comon}(\mathcal{V})$ is established.

Another piece of structure inherited from $\mathcal{V}$ to $\mathbf{Comon}(\mathcal{V})$ in the locally presentable context is monoidal closedness, again obtained from Theorem 2.12 for obtaining an adjoint for $- \otimes C \colon \mathbf{Comon}(\mathcal{V}) \to \mathbf{Comon}(\mathcal{V})$.

Proposition 2.15. [Por08b, 3.2] *If $\mathcal{V}$ is a locally presentable braided monoidal closed category, $\mathbf{Comon}(\mathcal{V})$ is also monoidal closed.*

2.3. **Universal measuring comonoid.** One of the basic goals of [HLFV15] was to estabilsh an enrichment of the category of monoids in the category of comonoids, under certain assumptions on $\mathcal{V}$. Below we summarize certain results; details can be found in Sections 4 and 5 therein. A basic tool is that of actions of monoidal categories.

Recall [JK02] that an *action* of a monoidal category on an ordinary one is given by a functor $* \colon \mathcal{V} \times \mathcal{D} \to \mathcal{D}$ expressing that $\mathcal{D}$ is a pseudomodule for the pseudomonoid $\mathcal{V}$ in the monoidal 2-category $(\mathbf{Cat}, \times, \mathbf{1})$. In more detail, we have natural isomorphisms with components

$$\chi_{X,Y,D} \colon (X \otimes Y) * D \xrightarrow{\sim} X * (Y * D) \quad \text{and} \quad \nu_D \colon I * D \xrightarrow{\sim} D \tag{7}$$

satisfying usual compatibility conditions. If $*$ is an action, it follows that $*^{\mathrm{op}}$ is an action too.

As a central example for our purposes, we have the action of the opposite monoidal category on itself via the internal hom, see [HLFV15, 3.7&5.1].

Lemma 2.16. *Suppose $\mathcal{V}$ is a braided monoidal closed category. The internal hom $[-,-] : \mathcal{V}^{\mathrm{op}} \times \mathcal{V} \to \mathcal{V}$ constitutes an action of $\mathcal{V}^{\mathrm{op}}$ on $\mathcal{V}$. Moreover, the induced $\mathbf{Mon}[-,-] : \mathbf{Comon}(\mathcal{V})^{\mathrm{op}} \times \mathbf{Mon}(\mathcal{V}) \to \mathbf{Mon}(\mathcal{V})$ (5) is an action of the monoidal $\mathbf{Comon}(\mathcal{V})^{\mathrm{op}}$ on $\mathbf{Mon}(\mathcal{V})$. Similarly for their opposite functors.*

A very important fact is that given a category $\mathcal{D}$ with an action from a monoidal category $\mathcal{V}$ with a parametrized adjoint, we obtain a $\mathcal{V}$-enriched category. This follows from a much stronger result of [GP97] regarding categories

enriched in bicategories; details can be found in [JK02] and [Vas14, § 4.3]. The explicit definitions of tensored and cotensored enriched categories can be found in [Kel05, 3.7].

Theorem 2.17. *Suppose that $\mathcal{V}$ is a monoidal category which acts on a category $\mathcal{D}$ via a functor $* : \mathcal{V} \times \mathcal{D} \to \mathcal{D}$, such that $- * D$ has a right adjoint $F(D, -)$ for every $D \in \mathcal{D}$ with a natural isomorphism*

$$\mathcal{D}(X * D, E) \cong \mathcal{V}(X, F(D, E)).$$

Then we can enrich $\mathcal{D}$ in $\mathcal{V}$, in the sense that there is a $\mathcal{V}$-category $\underline{\mathcal{D}}$ with hom-objects $\underline{\mathcal{D}}(A, B) = F(A, B)$ and underlying category $\mathcal{D}$. Moreover, if $\mathcal{V}$ is monoidal closed, the enrichment is tensored*, with $X * D$ the tensor of $X \in \mathcal{V}$ and $D \in \mathcal{D}$. If $\mathcal{V}$ is furthermore symmetric, the enrichment is* cotensored *if $X * -$ has a right adjoint; finally, we can also enrich $\mathcal{D}^{\mathrm{op}}$ in $\mathcal{V}$.*

By Lemma 2.16, the internal hom induces specific actions. Establishing parametrized adjoints for them will induce the desired enrichment of monoids in comonoids.

Theorem 2.18. [HLFV15, Thm 4.1] *If $\mathcal{V}$ is a locally presentable braided monoidal closed category, the functor $\mathbf{Mon}[-, B]^{\mathrm{op}} \colon \mathbf{Comon}(\mathcal{V}) \to \mathbf{Mon}(\mathcal{V})^{\mathrm{op}}$ has a right adjoint $P(-, B)$, i.e. there is a natural isomorphism*

$$\mathbf{Mon}(\mathcal{V})(A, [C, B]) \cong \mathbf{Comon}(\mathcal{V})(C, P(A, B)).$$

The parametrized adjoint of the functor $\mathbf{Mon}[-, -]^{\mathrm{op}}$, namely

$$P \colon \mathbf{Mon}(\mathcal{V})^{\mathrm{op}} \times \mathbf{Mon}(\mathcal{V}) \to \mathbf{Comon}(\mathcal{V}) \tag{8}$$

is called the *Sweedler hom*, and $P(A, B)$ is called the *universal measuring comonoid*, generalizing Sweedler's measuring coalgebras of [Swe69, §VII] as well as the *finite dual* $P(A, I) = A^o$ of an algebra A.

Moreover, each $\mathbf{Mon}[C, -]^{\mathrm{op}}$ turns out to also have a right adjoint $(C \triangleright -)^{\mathrm{op}}$ [Vas14, §6.2], and the induced functor of two variables

$$\triangleright \colon \mathbf{Comon}(\mathcal{V}) \times \mathbf{Mon}(\mathcal{V}) \to \mathbf{Mon}(\mathcal{V}) \tag{9}$$

is called the *Sweedler product* in [AJ13].

Applying Theorem 2.17 for the locally presentable braided monoidal closed category $\mathcal{V} = \mathbf{Comon}(\mathcal{V})$ by Prositions 2.13 and 2.15, and for $\mathcal{D} = \mathbf{Mon}(\mathcal{V})^{\mathrm{op}}$, we obtain the following part of [HLFV15, Thm. 5.2].

Theorem 2.19. *Suppose $\mathcal{V}$ is a locally presentable braided monoidal closed category.*

1. *The category $\mathbf{Mon}(\mathcal{V})^{\mathrm{op}}$ is a monoidal $\mathbf{Comon}(\mathcal{V})$-category, tensored and cotensored, with hom-objects $\underline{\mathbf{Mon}(\mathcal{V})^{\mathrm{op}}}(A, B) = P(B, A)$.*
2. *The category $\mathbf{Mon}(\mathcal{V})$ is a monoidal $\mathbf{Comon}(\mathcal{V})$-category, tensored and cotensored, with $\underline{\mathbf{Mon}(\mathcal{V})}(A, B) = P(A, B)$, cotensor $[C, B]$ and tensor $C \triangleright B$ for any comonoid C and monoid B.*

2.4. **Fibrations.** We recall some basic facts and constructions for the theory of fibrations and opfibrations, which is going to be of use in the paper. A few indicative references for the general theory are [Her94, Bor94b, Jac99], and for our specific context [Vas14].

Briefly recall that a functor $P\colon \mathcal{A} \to \mathbb{X}$ is a *fibration* when for every arrow $f\colon X \to Y$ in the *base* category $\mathbb{X}$ and every object B in the *total* category $\mathcal{A}$ above Y, i.e. $P(B) = Y$, there exists a *cartesian morphism* with codomain B above f. If we denote it $\phi\colon A \to B$, this means that for any $g\colon X \to X'$ and $A' \to B$ as in the diagram below, there exists a unique factorization through the domain of the cartesian morphism over g:



For each X in the base, its *fibre* category $\mathcal{A}_X \subset \mathcal{A}$ consists of objects above X and morphisms above the identity id_X, called *P-vertical*. We call ϕ a *cartesian lifting* of B along f. Assuming the axiom of choice, cartesian liftings can be selected (up to vertical isomorphism) for each morphism f in the base and object $B \in \mathcal{A}_{\mathrm{cod}f}$, henceforth denoted $\mathrm{Cart}(f, B)\colon f^*B \to B$.

Dually, a functor $U\colon \mathcal{C} \to \mathbb{X}$ is an *opfibration* when its opposite functor U^{op} is a fibration: for every $g\colon X \to Y$ in the base and $C \in \mathcal{C}_X$ above the domain, there is a cocartesian morphism from C above g, the *cocartesian lifting* of C along g denoted $\mathrm{Cocart}(g, C)\colon C \to g_!C$.

Any arrow in the total category of an (op)fibration factorizes uniquely into a vertical morphism followed by a (co)cartesian one:



(10)

The choice of (co)cartesian liftings in an (op)fibration induces a so-called *reindexing functor* between the fibre categories

$$f^* : \mathcal{A}_Y \to \mathcal{A}_X \quad \text{and} \quad g_!\colon \mathcal{C}_X \to \mathcal{C}_Y$$

respectively, for each morphism f or $g\colon X \to Y$ in the base category, mapping each object to the (co)domain of its lifting.

Remark 2.20. Due to the unique factorization of arrows in a (op)fibration through (co)cartesian liftings, we can deduce that a fibration $P : \mathcal{A} \to \mathbb{X}$ is also an opfibration (consequently a *bifibration*) if and only if, for every $f : X \to Y$ the reindexing $f^* : \mathcal{A}_Y \to \mathcal{A}_X$ has a left adjoint, namely $f_! : \mathcal{A}_X \to \mathcal{A}_Y$ (e.g. [Her93, Proposition 1.2.7]).

An *oplax fibred 1-cell* (S, F) between $P : \mathcal{A} \to \mathbb{X}$ and $Q : \mathcal{B} \to \mathbb{Y}$ is given by a commutative square of categories and functors



called an *oplax morphism of fibrations* in [Shu08, Def. 3.5]; this name is justified by the correspondence of Theorem 2.21.

By (10), in this case we always have a comparison vertical morphism



to the chosen Q-cartesian lifting of SB along Ff. If moreover S preserves cartesian arrows, meaning that if ϕ is P-cartesian then $S\phi$ is Q-cartesian or equivalently the above comparison map is an isomorphism, the pair (S, F) is called a *fibred 1-cell* or *strong morphism of fibrations*.

In particular, when P and Q are fibrations over the same base $\mathbb{X}$, we may consider oplax morphisms or fibred 1-cells of the form $(S, 1_{\mathbb{X}})$ displayed as



when S is called *(oplax) fibred functor*. Dually, we have the notion of an *lax opfibred 1-cell* (K, F), *opfibred 1-cell* when K is cocartesian, and *(lax) opfibred functor* $(K, 1_{\mathbb{X}})$.

Notice that any oplax fibred 1-cell (S, F) determines a collection of functors

$$S_X \colon \mathcal{A}_X \longrightarrow \mathcal{B}_{FX}$$

between the fibres, as the restriction of S to the corresponding subcategories.

An *fibred 2-cell* between oplax fibred 1-cells (S, F) and (T, G) is a pair of natural transformations $(\alpha : S \Rightarrow T, \beta : F \Rightarrow G)$ with α above β, *i.e.* $Q(\alpha_A) = \beta_{PA}$ for all $A \in \mathcal{A}$, displayed



A *fibred natural transformation* is of the form $(\alpha, 1_{1_{\mathbb{X}}}) : (S, 1_{\mathbb{X}}) \Rightarrow (T, 1_{\mathbb{X}})$ which ends up having vertical components, $Q(\alpha_A) = 1_{PA}$. Notice that if the 1-cells are strong, the definition of a 2-cell between them remains the same. Dually, we have the notion of an *opfibred 2-cell* and *opfibred natural transformation* between lax opfibred 1-cells and functors respectively.

We obtain 2-categories $\mathbf{Fib}_{\text{opl}}$ and $\mathbf{Fib}$ of fibrations over arbitrary base categories, (oplax) fibred 1-cells and fibred 2-cells. In particular, there are 2-categories $\mathbf{Fib}_{\text{opl}}(\mathbb{X})$ and $\mathbf{Fib}(\mathbb{X})$ of fibrations over a fixed base category $\mathbb{X}$, (oplax) fibred functors and fibred natural transformations. Dually, we have the 2-categories $\mathbf{OpFib}_{\text{(opl)}}$ and $\mathbf{OpFib}_{\text{(opl)}}(\mathbb{X})$.

The fundamental *Grothendieck construction* establishes a well-known equivalence between fibrations and pseudofunctors of Definition 2.2. Starting with a pseudofunctor $\mathscr{M} \colon \mathbb{X}^{\text{op}} \to \mathbf{Cat}$, we can form the Grothendieck category $\int \mathscr{M}$ with objects pairs $(A, X) \in \mathscr{M}X \times \mathbb{X}$ and morphisms $(A, X) \to (B, Y)$ pairs $(\phi \colon A \to (\mathscr{M}f)B, f \colon X \to Y) \in \mathscr{M}X \times \mathbb{X}$. This is a fibration over $\mathbb{X}$, with fibres $(\int \mathscr{M})_X = \mathscr{M}X$, reindexing functors $\mathscr{M}f$ and chosen cartesian liftings

$$\begin{array}{ccc} ((\mathscr{M}f)B, X) & \xrightarrow{(1_{(\mathscr{M}f)B}, f)} & (B, Y) \\ \vdots & & \vdots \\ X & \xrightarrow{\quad f \quad} & Y \end{array} \qquad \begin{array}{l} \text{in } \int \mathscr{M} \\ \\ \text{in } \mathbb{X}. \end{array} \tag{11}$$

Using similar machinery [Shu08, Prop. 3.6], we obtain respective correspondences for oplax fibred 1-cells and oplax natural transformations.

Theorem 2.21. *There are equivalences of 2-categories*

$$\mathbf{Fib}_{\text{opl}}(\mathbb{X}) \simeq [\mathbb{X}^{\text{op}}, \mathbf{Cat}]_{\text{opl}}$$
$$\mathbf{Fib}(\mathbb{X}) \simeq [\mathbb{X}^{\text{op}}, \mathbf{Cat}]$$

between the 2-categories of fibrations with fixed base and pseudofunctors with oplax or pseudonatural transformations and modifications.

There is also a 2-equivalence $\mathbf{ICat} \simeq \mathbf{Fib}$ between fibrations over arbitrary bases and an appropriately defined 2-category of pseudofunctors with arbitrary domain; for more details, see [Her93]. Along with the dual versions for opfibrations, namely $\mathbf{Fib}(\mathbb{X}) \simeq [\mathbb{X}, \mathbf{Cat}]$, these equivalences allow us to freely change our perspective between (op)fibrations and pseudofunctors.

Moving on to notions of adjunctions between fibrations, we obtain the following definitions as adjunctions in the respective 2-categories of fibrations.

Definition 2.22. Given fibrations $P : \mathcal{A} \to \mathbb{X}$ and $Q : \mathcal{B} \to \mathbb{Y}$, a *general (oplax) fibred adjunction* is given by a pair of (oplax) fibred 1-cells $(L, F) : P \to Q$ and $(R, G) : Q \to P$ together with fibred 2-cells $(\zeta, \eta) : (1_{\mathcal{A}}, 1_{\mathbb{X}}) \Rightarrow (RL, GF)$ and $(\xi, \varepsilon) : (LR, FG) \Rightarrow (1_{\mathcal{B}}, 1_{\mathbb{Y}})$ such that $L \dashv R$ via ζ, ξ and $F \dashv G$ via η, ε. This is displayed as

$$\begin{array}{ccc} \mathcal{A} & \underset{R}{\overset{L}{\rightleftarrows}} & \mathcal{B} \\ {\scriptstyle P}\downarrow & & \downarrow{\scriptstyle Q} \\ \mathbb{X} & \underset{G}{\overset{F}{\rightleftarrows}} & \mathbb{Y} \end{array}$$

and we write $(L, F) \dashv (R, G) : Q \to P$.

Notice that by definition, ζ is above η and ξ is above ε, hence (P, Q) is in particular an ordinary map between adjunctions. Dually, we have the notions of *general (lax) opfibred adjunction* and *opfibred adjunction* in $\mathbf{OpFib}_{(l)}$.

In [HLFV17, §3.2], conditions under which a fibred 1-cell has an adjoint are investigated in detail. Below we only recall the case of general oplax fibred adjunction, spelled out for opfibrations due to the applications that follow.

Theorem 2.23. *Suppose $(K, F) : U \to V$ is an opfibred 1-cell and $F \dashv G$ is an adjunction with counit ε between the bases of the opfibrations, as in*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{K} & \mathcal{D} \\ {\scriptstyle U}\downarrow & & \downarrow{\scriptstyle V} \\ \mathbb{X} & \underset{G}{\overset{F}{\underset{\longleftarrow}{\overset{\longrightarrow}{\bot}}}} & \mathbb{Y}. \end{array}$$

If, for each $Y \in \mathbb{Y}$, the composite functor between the fibres

$$\mathcal{C}_{GY} \xrightarrow{K_{GY}} \mathcal{D}_{FGY} \xrightarrow{(\varepsilon_Y)_!} \mathcal{D}_Y \tag{12}$$

has a right adjoint for each $Y \in \mathbb{Y}$, then K has a right adjoint R between the total categories and $(K, F) \dashv (R, G)$ is a general lax adjunction.

Finally, in [Vas17] the notion of an enriched fibration is discussed in length, and is closely related to [Vas14, §8.1] but with a relaxed condition which allows more examples to fit in the picture. Here, we gather the basic definitions in order to connect them with the setting of double categories later.

The following generalizes the notion of a (right) parametrized adjunction from **Cat** to $\mathbf{Fib}_{\mathrm{opl}}$ or $\mathbf{OpFib}_{\mathrm{l}}$.

Definition 2.24. A *general oplax parametrized adjunction* consists of oplax fibred 1-cells

$$\begin{array}{ccc} \mathcal{A} \times \mathcal{B} & \xrightarrow{F} & \mathcal{C} \\ {\scriptstyle H \times J}\downarrow & & \downarrow{\scriptstyle K} \\ \mathbb{X} \times \mathbb{Y} & \xrightarrow[G]{} & \mathbb{Z}, \end{array} \qquad \begin{array}{ccc} \mathcal{B}^{\mathrm{op}} \times \mathcal{C} & \xrightarrow{R} & \mathcal{A} \\ {\scriptstyle J^{\mathrm{op}} \times K}\downarrow & & \downarrow{\scriptstyle H} \\ \mathbb{Y}^{\mathrm{op}} \times \mathbb{Z} & \xrightarrow[S]{} & \mathbb{X} \end{array}$$

such the following is a general oplax fibred adjunction

$$\begin{array}{ccc} \mathcal{A} & \underset{R(B,-)}{\overset{F(-,B)}{\underset{\longleftarrow}{\overset{\longrightarrow}{\bot}}}} & \mathcal{C} \\ {\scriptstyle H}\downarrow & & \downarrow{\scriptstyle K} \\ \mathbb{X} & \underset{S(JB,-)}{\overset{G(-,JB)}{\underset{\longleftarrow}{\overset{\longrightarrow}{\bot}}}} & \mathbb{Z}. \end{array}$$

Dually, a *general lax parametrized adjunction* consists of lax opfibred 1-cells inducing a general lax opfibred adjunction as above.

Identifying the pseudomonoids in the cartesian monoidal 2-category **Fib**, we obtain the following definition which coincides with [Shu08, 12.1].

Definition 2.25. A fibration $T\colon \mathcal{V} \to \mathbb{W}$ is *monoidal* when $\mathcal{V}$, $\mathbb{W}$ are monoidal categories, T is a strict monoidal functor and the tensor product $\otimes_{\mathcal{V}}$ preserves cartesian arrows.

A fibration is *symmetric monoidal* when it is a strict braided monoidal functor between symmetric monoidal categories. Next, expressing a pseudomodule in $(\mathbf{Fib}, \times, \mathbf{1})$ gives the following.

Definition 2.26. A monoidal fibration $T : \mathcal{V} \to \mathbb{W}$ *acts* on the fibration $P : \mathcal{A} \to \mathbb{X}$ when there are two ordinary actions

$$\mu^{\mathcal{A}} = * : \mathcal{V} \times \mathcal{A} \longrightarrow \mathcal{A}$$
$$\mu^{\mathbb{X}} = \diamond : \mathbb{W} \times \mathbb{X} \longrightarrow \mathbb{X}$$

of the monoidal categories $\mathcal{V}$, $\mathbb{W}$ on the categories $\mathcal{A}$ and $\mathbb{X}$ respectively, such that $\mu^{\mathcal{A}}$ preserves cartesian arrows and the action constraints are above each other, in the sense that

$$P\chi^{\mathcal{A}}_{XYA} = \chi^{\mathbb{X}}_{(TX)(TY)(PA)}, \quad P\nu^{\mathcal{A}}_{A} = \nu^{\mathbb{X}}_{PA} \tag{13}$$

for all $X, Y \in \mathcal{V}$ and $A \in \mathcal{A}$, following notation (7).

Dually, we can define a *coaction* of a monoidal opfibration on an ordinary opfibration.

With the purpose of generalizing the establishment of an enrichment via actions from **Cat** in Theorem 2.17 to $\mathbf{Fib}_{\mathrm{opl}}$, we conclude to the following definition (see also [Vas14, 8.1.4]).

Definition 2.27. If $T\colon \mathcal{V} \to \mathbb{W}$ is a monoidal fibration, we say an ordinary fibration $P\colon \mathcal{A} \to \mathbb{X}$ is *enriched* in T when

- $\mathcal{A}$ is enriched in $\mathcal{V}$, $\mathbb{X}$ is enriched in $\mathbb{W}$ and the following commutes:

$$\begin{array}{ccc} \mathcal{A}^{\mathrm{op}} \times \mathcal{A} & \xrightarrow{\mathcal{A}(-,-)} & \mathcal{V} \\ {\scriptstyle P^{\mathrm{op}}\times P}\downarrow & & \downarrow{\scriptstyle T} \\ \mathbb{X}^{\mathrm{op}} \times \mathbb{X} & \xrightarrow[\mathbb{X}(-,-)]{} & \mathbb{W} \end{array}$$

- composition and identities of the enrichments are compatible, in that

$$TM^{\mathcal{A}}_{A,B,C} = M^{\mathbb{X}}_{PA,PB,PC} \quad \text{and} \quad Tj^{\mathcal{A}}_{A} = j^{\mathbb{X}}_{PA}.$$

Dually, we have the notion of an opfibration enriched in a monoidal opfibration, and we also introduce the following definition.

Definition 2.28. Suppose that $T : \mathcal{V} \to \mathbb{W}$ is a symmetric monoidal opfibration. We say that a fibration $P : \mathcal{A} \to \mathbb{X}$ is enriched in T if the opfibration $P^{\mathrm{op}} : \mathcal{A}^{\mathrm{op}} \to \mathbb{X}^{\mathrm{op}}$ is an enriched T-opfibration.

Finally, the following theorem gives a direct way of obtaining an enriched fibration from actions both between the total and the base categories, and is part of a stronger argument in [Vas14, 8.1.6].

Theorem 2.29. *Suppose that $T : \mathcal{V} \to \mathbb{W}$ is a monoidal fibration, which acts on an (ordinary) fibration $P : \mathcal{A} \to \mathbb{X}$ via the fibred 1-cell*

$$\begin{array}{ccc} \mathcal{V} \times \mathcal{A} & \xrightarrow{\ *\ } & \mathcal{A} \\ {\scriptstyle T\times P}\downarrow & & \downarrow{\scriptstyle P} \\ \mathbb{W} \times \mathbb{X} & \xrightarrow[\diamond]{} & \mathbb{X}. \end{array}$$

If this action has an oplax fibred parametrized adjoint $(R, S) : P^{\mathrm{op}} \times P \to T$, then we can enrich the fibration P in the monoidal fibration T.

We will later use the dual version, for which an action of a monoidal opfibration induces an enrichment via a general lax opfibred parametrized adjunction.

3. Double categories

The setting of double categories, and more specifically fibrant monoidal double categories, is crucial for this work's development but also for further possible applications. A few references for the theory of double categories and results relevant here are [GP99, GP04, Shu08]; the original concept of a double category as a category internal in **Cat** goes back to [Ehr63]. In order to provide a passage from double categories to bicategories, we largely follow the notation and approach of [Shu10] where a method for constructing monoidal bicategories from monoidal double categories is described.

We then proceed to the study of monads (see also [FGK11]) and comonads in double categories, as well a natural (op)fibrational picture they form over the vertical category induced by fibrancy conditions. In the monoidal case, these are in fact monoidal (op)fibrations in the sense of the previous section.

Finally, by introducing the notion of a *locally monoidal closed* double category, capturing a closed structure for both vertical and horizontal categories, we are able to explore enrichment relations between the category of monads and comonads, sometimes forming an enriched fibration.

3.1. Background on fibrant double categories.

We recall the central definitions and fix our notation.

Definition 3.1. A *(pseudo) double category* $\mathbb{D}$ consists of a category of objects $\mathbb{D}_0$ and a category of arrows $\mathbb{D}_1$, with (identity, source and target, composition) structure functors

$$\mathbf{1} : \mathbb{D}_0 \to \mathbb{D}_1, \quad \mathfrak{s}, \mathfrak{t} : \mathbb{D}_1 \rightrightarrows \mathbb{D}_0, \quad \odot : \mathbb{D}_1 \times_{\mathbb{D}_0} \mathbb{D}_1 \to \mathbb{D}_1$$

such that $\mathfrak{s}(1_A)=\mathfrak{t}(1_A)=A$, $\mathfrak{s}(M \odot N)=\mathfrak{s}(N)$, $\mathfrak{t}(M \odot N)=\mathfrak{t}(M)$ for all $A \in \mathrm{ob}\mathbb{D}_0$ and $M, N \in \mathrm{ob}\mathbb{D}_1$, equipped with natural isomorphisms

$$\alpha : (M \odot N) \odot P \xrightarrow{\sim} M \odot (N \odot P)$$
$$\lambda : 1_{\mathfrak{s}(M)} \odot M \xrightarrow{\sim} M \quad \rho : M \odot 1_{\mathfrak{t}(M)} \xrightarrow{\sim} M$$

in $\mathbb{D}_1$ such that $\mathfrak{t}(\alpha), \mathfrak{s}(\alpha), \mathfrak{t}(\lambda), \mathfrak{s}(\lambda), \mathfrak{t}(\rho), \mathfrak{s}(\rho)$ are all identities, and satisfying the usual coherence conditions (as for a bicategory (1)).

The objects of $\mathbb{D}_0$ are called *0-cells* and the morphisms $f : A \to B$ of $\mathbb{D}_0$ are called *vertical 1-cells*. The objects of $\mathbb{D}_1$ are the *horizontal 1-cells* or *proarrows*, $M : A \nrightarrow B$. The morphisms of $\mathbb{D}_1$ are the *2-morphisms*

$$\begin{array}{ccc} A & \overset{M}{\nrightarrow} & B \\ {\scriptstyle f}\downarrow & \Downarrow\alpha & \downarrow{\scriptstyle g} \\ C & \underset{N}{\nrightarrow} & D \end{array} \tag{14}$$

or ${}^{f}\alpha^{g} : M \Rightarrow N$, where $\mathfrak{s}(\alpha) = f$ and $\mathfrak{t}(\alpha) = g$. The composition of vertical 1-cells and the vertical composition of 2-morphisms are strictly associative, whereas horizontal composition of horizontal 1-cells and 2-morphisms is associative up to isomorphism, written

$$\begin{array}{ccc} A & \overset{M}{\nrightarrow} & B \\ {\scriptstyle f}\downarrow & \Downarrow\alpha & \downarrow{\scriptstyle g} \\ C & \overset{N}{\nrightarrow} & D \\ {\scriptstyle h}\downarrow & \Downarrow\beta & \downarrow{\scriptstyle k} \\ E & \underset{P}{\nrightarrow} & F \end{array} = \begin{array}{ccc} A & \overset{M}{\nrightarrow} & B \\ {\scriptstyle hf}\downarrow & \Downarrow\beta\alpha & \downarrow{\scriptstyle kg} \\ E & \underset{P}{\nrightarrow} & F, \end{array} \qquad \begin{array}{ccccc} A & \overset{M}{\nrightarrow} & B & \overset{N}{\nrightarrow} & C \\ {\scriptstyle f}\downarrow & \Downarrow\alpha g & \downarrow & \Downarrow\beta & \downarrow{\scriptstyle h} \\ D & & \underset{P}{\nrightarrow} & E & \underset{K}{\nrightarrow} F \end{array} = \begin{array}{ccc} A & \overset{N\odot M}{\nrightarrow} & C \\ {\scriptstyle f}\downarrow & \Downarrow\beta\odot\alpha & \downarrow{\scriptstyle h} \\ D & \underset{K\odot P}{\nrightarrow} & F. \end{array} \tag{15}$$

Strict identities are $\mathrm{id}_A : A \to A$ and ${}^{\mathrm{id}_A}1_M^{\mathrm{id}_B} : M \Rightarrow M$, and horizontal units are $1_A : A \nrightarrow A$ and ${}^{f}1_f^{f} : 1_A \Rightarrow 1_B$. Functoriality of the horizontal composition results in the relation $1_N \odot 1_M = 1_{N\odot M}$ and the interchange law $(\beta'\beta)\odot(\alpha'\alpha) = (\beta'\odot\alpha')(\beta\odot\alpha)$. A 2-morphism with identity source and target vertical 1-cell, like a, l, r above, is called *globular*.

The *opposite double category* $\mathbb{D}^{\mathrm{op}}$ is the double category with vertical category $\mathbb{D}_0^{\mathrm{op}}$ and horizontal category $\mathbb{D}_1^{\mathrm{op}}$. There also exist the *horizontally opposite* double category $\mathbb{D}^{\mathrm{hop}}$ and *vertically opposite* double category $\mathbb{D}^{\mathrm{vop}}$ with opposite horizontal and vertical categories respectively.

For every double category $\mathbb{D}$ there is a corresponding bicategory denoted by $\mathcal{H}(\mathbb{D})$ called its *horizontal bicategory*. It consists of the objects, horizontal 1-cells and globular 2-morphisms, and clearly are required conditions are satisfied. In a sense, this comes from discarding the vertical structure of the double category. Many well-known bicategories arise as the horizontal bicategories of specific double categories; in fact, all the examples of Section 2.1 can be seen as the horizontal ones for the following double categories, see e.g. [GP99, Shu08], with appropriate 2-morphisms.

- $\mathbb{S}\mathbf{pan}(\mathcal{C})$ and $\mathbb{R}\mathbf{el}(\mathcal{C})$ with vertical category $\mathcal{C}$;
- $\mathbb{B}\mathbf{Mod}$ with vertical category $\mathbf{Rng}$ of rings and ring homomorphisms;
- $(\mathcal{V}\text{-})\mathbb{P}\mathbf{rof}$ with vertical category $(\mathcal{V}\text{-})\mathbf{Cat}$.

What is evident from the above examples is that the double categorical perspective includes not only the morphisms which the bicategorical structure is named after and describes best, but also the more fundamental, strict morphisms between the objects: functions, ring maps and functors above.

Definition 3.2. For $\mathbb{D}$ and $\mathbb{E}$ (pseudo) double categories, a *pseudo double functor* $F : \mathbb{D} \to \mathbb{E}$ consists of functors $F_0 : \mathbb{D}_0 \to \mathbb{E}_0$ and $F_1 : \mathbb{D}_1 \to \mathbb{E}_1$ between the categories of objects and arrows, such that $\mathfrak{s} \circ F_1 = F_0 \circ \mathfrak{s}$ and

$\mathfrak{t} \circ F_1 = F_0 \circ \mathfrak{t}$, and natural transformations $F_\odot$, F_U with components globular isomorphisms $F_1 M \odot F_1 N \xrightarrow{\sim} F_1(M \odot N)$ and $1_{F_0 A} \xrightarrow{\sim} F_1(1_A)$ which satisfy the usual coherence axioms (2) and (3) for a pseudofunctor.

Due to the compatibility of F_0, F_1 with sources and targets, we can write the mapping of F_1 on 1-cells and 2-morphisms as

$$\begin{array}{ccc} A & \xrightarrow{M} & B \\ {\scriptstyle f}\downarrow & \Downarrow\alpha & \downarrow{\scriptstyle g} \\ C & \xrightarrow[N]{} & D \end{array} \quad \mapsto \quad \begin{array}{ccc} F_0 A & \xrightarrow{F_1 M} & F_0 B \\ {\scriptstyle F_0 f}\downarrow & \Downarrow F_1\alpha & \downarrow{\scriptstyle F_0 g} \\ F_0 C & \xrightarrow[F_1 N]{} & F_0 D \end{array} \tag{16}$$

We also have notions of *lax* and *colax double functors* between pseudo double categories, where the natural transformations $F_\odot$ and F_U have components globular 2-morphisms in one of the two possible directions respectively. The full definitions can be found in the appendix of [GP99] or [GP04].

Any lax/colax/pseudo double functor $F : \mathbb{D} \to \mathbb{E}$ naturally induces a lax/colax/pseudo functor between the respective horizontal bicategories, $\mathcal{H}F : \mathcal{H}(\mathbb{D}) \to \mathcal{H}(\mathbb{E})$, where each $A \in \mathbb{D}_0$ is mapped to $F_0 A \in \mathbb{E}_0$, and there are ordinary functors $\mathcal{H}F_{A,B} : \mathcal{H}(\mathbb{D})(A, B) \to \mathcal{H}(\mathbb{E})(F_0 A, F_0 B)$ mapping globular 2-cells to globular 2-cells via (16).

With an appropriate notion for transformations between double functors, there is a 2-category $\mathcal{D}bl$ of double categories. We can then obtain the definition of a monoidal double category as a pseudomonoid therein, see [Shu10, 2.9] for the full definition. Notably, in [GP04] the tensor product $\otimes$ is required to be a colax double functor rather than pseudo double.

Definition 3.3. A *monoidal double category* is a double category $\mathbb{D}$ equipped with pseudo double functors $\otimes = (\otimes_1, \otimes_1) : \mathbb{D} \times \mathbb{D} \to \mathbb{D}$ and $\mathbf{I} : \mathbf{1} \to \mathbb{D}$ and invertible transformations expressing associativity and unity constraints, subject to usual axioms.

This amounts to $(\mathbb{D}_0, \otimes_0, I)$ and $(\mathbb{D}_1, \otimes_1, 1_I)$ being monoidal categories with $I = \mathbf{I}(*)$ and $1_I : I \nrightarrow I$, the functors $\mathfrak{s}, \mathfrak{t}$ being strict monoidal and preserving associativity and unit constraints, and globular isomorphisms

$$\begin{gathered} (M \otimes_1 N) \odot (M' \otimes_1 N') \cong (M \odot M') \otimes_1 (N \odot N') \\ 1_{(A \otimes_0 B)} \cong 1_A \otimes_1 1_B \end{gathered} \tag{17}$$

subject to coherence conditions.

A *braided* or *symmetric* monoidal double category $\mathbb{D}$ is one for which $\mathbb{D}_0$, $\mathbb{D}_1$ are braided or symmetric, and the source and target functors $\mathfrak{s}, \mathfrak{t}$ are strict braided monoidal, subject subject to two more axioms.

By definition (16), $\otimes_1$ is the following mapping:

$$\otimes_1 : \mathbb{D}_1 \times \mathbb{D}_1 \longrightarrow \mathbb{D}_1 \tag{18}$$

$$\begin{array}{ccc} (A & \xrightarrow{M} & B, \quad C \xrightarrow{N} D) \\ {\scriptstyle f}\downarrow & \Downarrow\alpha & \downarrow{\scriptstyle g} \quad {\scriptstyle h}\downarrow \quad \Downarrow\beta \quad \downarrow{\scriptstyle k} \\ (A' & \xrightarrow[M']{} & B', \quad C' \xrightarrow[N']{} D') \end{array} \mapsto \begin{array}{ccc} A \otimes_0 C & \xrightarrow{M \otimes_1 N} & B \otimes_0 D \\ {\scriptstyle f \otimes_0 h}\downarrow & \Downarrow\alpha \otimes_1 \beta & \downarrow{\scriptstyle g \otimes_0 k} \\ A' \otimes_0 B' & \xrightarrow[M' \otimes_1 N']{} & B' \otimes_1 D' \end{array}$$

Passing on to the theory of fibrant double categories, it is the case that in many important examples there exists a canonical way of turning vertical 1-cells into horizontal 1-cells. Such links have been studied in various works, and the terminology used below can be found in [GP04, Shu08, DPP10].

Definition 3.4. Let $\mathbb{D}$ be a double category and $f : A \to B$ a vertical 1-cell. A *companion* of f is a horizontal 1-cell $\hat{f} : A \nrightarrow B$ together with 2-morphisms



such that $p_1 p_2 = 1_f$ and $p_1 \odot p_2 \cong 1_{\hat{f}}$. Dually, a *conjoint* of f is a horizontal 1-cell $\check{f} : B \nrightarrow A$ together with 2-morphisms



such that $q_1 q_2 = 1_f$ and $q_2 \odot q_1 \cong 1_{\check{f}}$.

The ideas which led to the above definitions go back to [BS76], where a *connection* on a double category corresponds to a strictly functorial choice of a companion for each vertical arrow.

Definition 3.5. [Shu10, Definition 3.4] A *fibrant double category* is a double category for which every vertical 1-morphism has a companion and a conjoint.

Fibrant double categories are also called *framed bicategories* or equivalently *proarrow equipments* [Woo82, Ver11]. The above definition's equivalence with the following can be found for example at [Shu08, Thm. 4.1].

Definition 3.6. A fibrant double category is one where the functor

$$(\mathfrak{s}, \mathfrak{t}) \colon \mathbb{D}_1 \longrightarrow \mathbb{D}_0 \times \mathbb{D}_0$$

mapping each horizontal 1-cell and 2-morphism to the pair of source and target is a fibration, or equivalently an opfibration.

In this view, the canonical cartesian lifting of some $N \colon C \nrightarrow D$ along a pair of vertical morphisms $f \colon A \to C$, $g \colon B \to D$



(19)

Many properties for fibrant double categories can be deduced from the companion and conjoint definitions. The following lemma gathers the most useful

for us; the explicit proofs can be found in [Shu10], or can be easily deduced e.g. via mates and factorization through lifting (10) for Item 5.

Lemma 3.7. *Suppose $\mathbb{D}$ is a fibrant double category.*

(1) Companions and conjoints of a vertical 1-cell are essentially unique (up to unique globular isomorphism).
(2) The horizontal composites $\hat{g} \odot \hat{f}$ and $\check{g} \odot \check{f}$ are the companion and the conjoint of the vertical composite gf.
(3) The companion and conjoint of the vertical identities are the horizontal identites, $\widehat{\mathrm{id}_A} = \widecheck{\mathrm{id}_A} = 1_A$.
(4) For any vertical 1-cell $f\colon A \to B$, we have an adjunction $\hat{f} \dashv \check{f}$ in the horizontal bicategory $\mathcal{H}(\mathbb{D})$.
(5) For any vertical 1-cells $f\colon A \to C$, $g\colon B \to D$ and horizontal 1-cells $M\colon A \nrightarrow B$, $N\colon C \nrightarrow D$, we have bijections between

$$\mathcal{H}(\mathbb{D})(M, \hat{f} \odot N \odot \check{g}) \cong \mathcal{H}(\mathbb{D})(M \odot \hat{g}, \hat{f} \odot N) \cong \mathcal{H}(\mathbb{D})(\check{f} \odot M, N \odot \check{g}) \cong \mathcal{H}(\mathbb{D})(\check{f} \odot M \odot \hat{g}, N) \tag{20}$$

and the 2-morphisms $M \Rightarrow N$ with source and target f and g (14).
(6) If $\mathbb{D}$ is also monoidal, $\hat{f} \otimes_1 \hat{g}$ and $\check{f} \otimes_1 \check{g}$ are the companion and conjoint of $f \otimes_0 g$ for any vertical 1-cells f, g.

For example, $\mathcal{V}$-$\mathbb{P}\mathbf{rof}$ is a fibrant double category: the companion and conjoint for a $\mathcal{V}$-functor $F\colon \mathcal{A} \to \mathcal{B}$ are given by the *representable profunctors*

$$\hat{F} = F_*\colon \mathcal{A} \nrightarrow \mathcal{B} \text{ by } F_*(B, A) = \mathcal{B}(B, FA)$$

$$\check{F} = F^*\colon \mathcal{B} \nrightarrow \mathcal{A} \text{ by } F^*(A, B) = \mathcal{B}(FA, B)$$

For $\mathbb{S}\mathbf{pan}(\mathcal{C})$, the companion and conjoint of a function $f\colon A \to B$ are $\check{f} = (\mathrm{id}_A, f)$ and $\hat{f} = (f, \mathrm{id}_B)$, whereas for $\mathbf{BMod}$ a ring morphism $f\colon A \to B$ gives rise to B as a left-A right-B bimodule but also as a left-B right-A bimodule via restriction of scalars.

A fundamental property of a fibrant double category with a monoidal structure is that its horizontal bicategory inherits it. This process, studied in detail in [Shu10], allows us to reduce a lengthy and demanding task of verifying the coherence conditions of monoidal structure on a bicategory into a much more concise one, essentially involving a pair of ordinary monoidal categories.

Theorem 3.8. [Shu10, Theorem 5.1] *If $\mathbb{D}$ is a fibrant monoidal double category, then $\mathcal{H}(\mathbb{D})$ is a monoidal bicategory. If $\mathbb{D}$ is braided or symmetric, then so is $\mathcal{H}(\mathbb{D})$.*

Evidently, the monoidal structure of the bicategory consists of the induced pseudofunctor of bicategories $\mathcal{H}(\otimes) : \mathcal{H}(\mathbb{D}) \times \mathcal{H}(\mathbb{D}) \to \mathcal{H}(\mathbb{D})$ and the monoidal unit 1_I of $\mathbb{D}_1$, for a $(\mathbb{D}, \otimes, I)$ as in Definition 3.3.

3.2. **Monads and comonads in double categories.** Suppose $\mathbb{D}$ is a double category. Define the category of *endomorphisms* $\mathbb{D}_1^\bullet$ to be the (non-full) subcategory of $\mathbb{D}_1$ of all horizontal endo-1-cells and 2-morphisms with the same

source and target. Explicitly, objects are of the form $M : A \nrightarrow A$ and arrows

$$\begin{array}{ccc} A & \overset{M}{\nrightarrow} & A \\ {\scriptstyle f}\downarrow & \Downarrow\alpha & \downarrow{\scriptstyle f} \\ B & \underset{N}{\nrightarrow} & B \end{array} \tag{21}$$

denoted by $\alpha_f : M_A \to N_B$. In [FGK11], this category coincides with the vertical 1-category of the double category $\mathbb{E}\mathbf{nd}(\mathbb{D})$ of (horizontal) endomorphisms, horizontal endomorphism maps, vertical endomorphism maps and endomorphism squares in $\mathbb{D}$.

Definition 3.9. A *monad* in a double category $\mathbb{D}$ is a horizontal endo-1-cell $M : A \nrightarrow A$ *i.e.* an object in $\mathbb{D}_1^\bullet$, equipped with globular 2-morphisms

$$\begin{array}{ccccc} A & \overset{M}{\nrightarrow} & A & \overset{M}{\nrightarrow} & A \\ {\scriptstyle \mathrm{id_A}}\downarrow & & \Downarrow m & & \downarrow{\scriptstyle \mathrm{id}_A} \\ A & & \underset{M}{\nrightarrow} & & A, \end{array} \qquad \begin{array}{ccc} A & \overset{1_A}{\nrightarrow} & A \\ {\scriptstyle \mathrm{id_A}}\downarrow & \Downarrow\eta & \downarrow{\scriptstyle \mathrm{id}_A} \\ A & \underset{M}{\nrightarrow} & A \end{array}$$

satisfying the usual associativity and unit laws; this is the same as a monad in its horizontal bicategory $\mathcal{H}(\mathbb{D})$ (Definition 2.5). A *monad homomorphism* consists of an arrow $\alpha_f : M_A \to N_B$ in $\mathbb{D}_1^\bullet$ which respects multiplication and unit:

$$\begin{array}{ccccc} A & \overset{M}{\nrightarrow} & A & \overset{M}{\nrightarrow} & A \\ {\scriptstyle f}\downarrow & \Downarrow\alpha & \downarrow{\scriptstyle f} & \Downarrow\alpha & \downarrow{\scriptstyle f} \\ B & \underset{N}{\nrightarrow} & B & \underset{N}{\nrightarrow} & B \\ \| & & \Downarrow m & & \| \\ B & & \underset{N}{\nrightarrow} & & B \end{array} = \begin{array}{ccccc} A & \overset{M}{\nrightarrow} & A & \overset{M}{\nrightarrow} & A \\ \| & & \Downarrow m & & \| \\ A & & \underset{M}{\nrightarrow} & & A \\ {\scriptstyle f}\downarrow & & \Downarrow\alpha & & \downarrow{\scriptstyle f} \\ B & & \underset{N}{\nrightarrow} & & B, \end{array} \qquad \begin{array}{ccc} A & \overset{1_A}{\nrightarrow} & A \\ \| & \Downarrow\eta & \| \\ A & \underset{M}{\nrightarrow} & A \\ {\scriptstyle f}\downarrow & \Downarrow\alpha & \downarrow{\scriptstyle f} \\ B & \underset{N}{\nrightarrow} & B \end{array} = \begin{array}{ccc} A & \overset{1_A}{\nrightarrow} & A \\ {\scriptstyle f}\downarrow & \Downarrow 1_f & \downarrow{\scriptstyle f} \\ B & \underset{1_B}{\nrightarrow} & B \\ \| & \Downarrow\eta & \| \\ B & \underset{N}{\nrightarrow} & B. \end{array} \tag{22}$$

We obtain a non-full subcategory of $\mathbb{D}_1$, the category $\mathbf{Mnd}(\mathbb{D})$. These definitions can be found in [Shu08] under the names of *monoids* and *monoid homomorphisms* for fibrant double categories, as well as in [FGK11] as monads and *vertical* monad maps in a double category $\mathbb{D}$. In the latter work, $\mathbf{Mnd}(\mathbb{D})$ is the vertical category of $\mathbb{M}\mathbf{nd}(\mathbb{D})$, a double category of monads, horizontal and vertical monad maps and monad squares.

Dually, we have the following definition.

Definition 3.10. There is a category $\mathbf{Cmd}(\mathbb{D})$ with objects *comonads* in $\mathbb{D}$, *i.e.* horizontal endo-1-cells $C : A \nrightarrow A$ equipped with globular 2-morphisms

$$\begin{array}{ccccc} A & & \overset{C}{\nrightarrow} & & A \\ {\scriptstyle \mathrm{id_A}}\downarrow & & \Downarrow\Delta & & \downarrow{\scriptstyle \mathrm{id}_A} \\ A & \underset{C}{\nrightarrow} & A & \underset{C}{\nrightarrow} & A, \end{array} \qquad \begin{array}{ccc} A & \overset{C}{\nrightarrow} & A \\ {\scriptstyle \mathrm{id_A}}\downarrow & \Downarrow\epsilon & \downarrow{\scriptstyle \mathrm{id}_A} \\ A & \underset{C}{\nrightarrow} & A \end{array}$$

satisfying the usual coassociativity and counit axioms for a comonad in the horizontal bicategory $\mathcal{H}(\mathbb{D})$ (Definition 2.8). Arrows are *comonad homomorphisms*, *i.e.* $\alpha_f : C_A \to D_B$ in $\mathbb{D}_1^\bullet$ satisfying dual axioms to (22).

Observe that $\mathbf{Mnd}(\mathbb{D}^{\mathrm{op}}) = \mathbf{Cmd}(\mathbb{D})^{\mathrm{op}}$, and that the forgetful functors $\mathbf{Mnd}(\mathbb{D}), \mathbf{Cmd}(\mathbb{D}) \to \mathbb{D}_1^\bullet \to \mathbb{D}_1$ reflect isomorphisms. Moreover, there are also forgetful functors to $\mathbb{D}_0$, mapping a horizontal endo-1-cell $M : A \nrightarrow A$ to its source and target A, and a 2-morphism $\alpha_f \colon M_A \to N_B$ to its source and target $f \colon A \to B$; these are studied in detail below.

Although in [Vas14, §8] the above structures were called (co)monoids, this notation is preferred due to the fact that it doesn't require any monoidal structure on the double category. A monoid in a monoidal double category should correspond to a lax double functor $\mathbf{1} \to \mathbb{D}$ which comes down to a monoid in the vertical category $\mathbb{D}_0$ as the source and target of a monoid in the horizontal category $\mathbb{D}_1$.

We now consider how different notions of double functors relate to the categories of endomorphisms, monads and comonads. The following can be deduced from the definition of a double functor (16) in a straightforward way.

Corollary 3.11. *Suppose that $F \colon \mathbb{D} \to \mathbb{E}$ is a lax/colax/pseudo double functor. Then F_1 naturally induces an ordinary functor $F_1^\bullet \colon \mathbb{D}_1^\bullet \to \mathbb{E}_1^\bullet$.*

For monads and comonads, the following resembles to standard properties of monoidal functors, see Section 2.2.

Proposition 3.12. *Any lax double functor $(F_0, F_1) : \mathbb{D} \to \mathbb{E}$ induces an ordinary functor*

$$\mathbf{Mon}F : \mathbf{Mnd}(\mathbb{D}) \to \mathbf{Mnd}(\mathbb{E})$$

between the categories of monads, given by restricting F_1 to $\mathbf{Mnd}(\mathbb{D})$. Dually, any colax double functor induces a functor between the categories of comonoids,

$$\mathbf{Comon}F : \mathbf{Cmd}(\mathbb{D}) \to \mathbf{Cmd}(\mathbb{E}).$$

Proof. A monad $M : A \nrightarrow A$ with $m : M \odot M \to M$ and $\eta : 1_M \to M$ is mapped to $F_1 M : F_0 A \nrightarrow F_0 A$ with multiplication and unit

$$\begin{array}{ccccc} F_0A & \xrightarrow{F_1M} & F_0A & \xrightarrow{F_1M} & F_0A \\ \| & & \Downarrow F_\odot & & \| \\ F_0A & & \xrightarrow[F_1(M\odot M)]{} & & F_0A \\ \| & & \Downarrow F_1 m & & \| \\ F_0A & & \xrightarrow[F_1M]{} & & F_0A \end{array} \qquad \text{and} \qquad \begin{array}{ccc} F_0A & \xrightarrow{F_1(1_A)} & F_0A \\ \| & \Downarrow F_U & \| \\ F_0A & \xrightarrow[1_{F_0A}]{} & F_0A \\ \| & \Downarrow F_1\eta & \| \\ F_0A & \xrightarrow[F_1M]{} & F_0A \end{array}$$

and the axioms follow from the axioms for $F_\odot$ and F_U. A monad homomorphism $\alpha_f : M_A \to N_B$ is mapped to $F_1^\bullet \alpha$,

$$\begin{array}{ccc} F_0A & \xrightarrow{F_1M} & F_0A \\ {\scriptstyle F_0f}\downarrow & \Downarrow F_1\alpha & \downarrow{\scriptstyle F_0f} \\ F_0B & \xrightarrow[F_1N]{} & F_0B \end{array}$$

which respects multiplications and units by naturality of $F_\odot$ and F_U. Similarly for the induced functor between comonads. □

Remark 3.13. Recall by Remark 2.9 that monads in a bicategory are monoids in a monoidal endohom category with horizontal composition. This point of view will be useful later, hence we should equivalently view double categorical monads as $M_A \in \mathbf{Mon}(\mathcal{H}(\mathbb{D})(A,A), \odot, 1_A)$ and comonads as $C_A \in \mathbf{Comon}(\mathcal{H}(\mathbb{D})(A,A), \odot, 1_A)$. (Co)monad homomorphisms cannot be expressed in a similar way, since the respective 2-morphisms do not have to be globular, but this is precisely the reason why categories of double (co)monads capture further desired structure.

Since (co)lax double functors induce (co)lax functors between the horizontal bicategories, on the level of objects Proposition 3.12 coincides with Remark 2.6.

As an application of the above, consider a monoidal double category $(\mathbb{D}, \otimes, \mathbf{I})$ as in Definition 3.3. The pseudo double functor $\otimes\colon \mathbb{D} \times \mathbb{D} \to \mathbb{D}$ induces, by Corollary 3.11 and Proposition 3.12, ordinary functors

$$\otimes_1^\bullet\colon \mathbb{D}_1^\bullet \times \mathbb{D}_1^\bullet \to \mathbb{D}_1^\bullet$$
$$\mathbf{Mon}\otimes\colon \mathbf{Mnd}(\mathbb{D}) \times \mathbf{Mnd}(\mathbb{D}) \to \mathbf{Mnd}(\mathbb{D})$$
$$\mathbf{Comon}\otimes\colon \mathbf{Cmd}(\mathbb{D}) \times \mathbf{Cmd}(\mathbb{D}) \to \mathbf{Cmd}(\mathbb{D})$$

given by $\otimes_1$ (18) restricted to the specific subcategories of $\mathbb{D}_1$. Along with the monoidal unit $1_I : I \nrightarrow I$, and since the forgetful functors to the monoidal $\mathbb{D}_1$ are conservative, we obtain the following.

Proposition 3.14. *If $\mathbb{D}$ is a monoidal double category, then the categories $\mathbb{D}_1^\bullet$, $\mathbf{Mnd}(\mathbb{D})$ and $\mathbf{Cmd}(\mathbb{D})$ inherit a monoidal structure from $\mathbb{D}_1$. When $\mathbb{D}$ is braided or symmetric, then so are the categories of endomorphisms, monads and comonads.*

We would now like to further study the categories of endomorphisms, monads and comonads in fibrand double categories Definition 3.5.

Proposition 3.15. *If $\mathbb{D}$ is a fibrant double category, $\mathbb{D}_1^\bullet$ is bifibred over $\mathbb{D}_0$.*

Proof. Due to the correspondence of Theorem 2.21, it is enough to define pseudofunctors from $\mathbb{D}_0^{(\mathrm{op})}$ which will give rise to a fibration and opfibration with total categories isomorphic to $\mathbb{D}_1^\bullet$ via the Grothendieck construction; define

$$\begin{array}{ccc} \mathscr{M}\ :\ \mathbb{D}_0^{\mathrm{op}} & \longrightarrow & \mathbf{Cat}, \\ A & \mapsto & \mathcal{H}(\mathbb{D})(A,A) \\ {\scriptstyle f}\downarrow & & \uparrow{\scriptstyle (\check{f}\odot\text{-}\odot\hat{f})} \\ B & \mapsto & \mathcal{H}(\mathbb{D})(B,B) \end{array} \qquad \begin{array}{ccc} \mathscr{F}\ :\ \mathbb{D}_0 & \longrightarrow & \mathbf{Cat} \\ A & \mapsto & \mathcal{H}(\mathbb{D})(A,A) \\ {\scriptstyle f}\downarrow & & \downarrow{\scriptstyle (\hat{f}\odot\text{-}\odot\check{f})} \\ B & \mapsto & \mathcal{H}(\mathbb{D})(B,B). \end{array} \tag{23}$$

In more detail, the first one is given by the mapping on objects and arrows

$$\left(B \;\overset{H}{\underset{H'}{\Downarrow\sigma}}\; B\right) \mapsto \left(A \xrightarrow{\hat{f}} B \;\overset{H}{\underset{H'}{\Downarrow\sigma}}\; B \xrightarrow{\check{f}} A\right)$$

pre-composing with the companion and post-composing with the conjoint of the given vertical 1-cell. For these mappings to constitute a pseudofunctor $\mathscr{M}$, we need certain natural isomorphisms satisfying coherence conditions as

in Definition 2.2. For every triple of 0-cells A, B, C, there is a natural isomorphism δ with components

$$\begin{array}{ccccc} & & \mathcal{H}(\mathbb{D})(B,B) & & \\ & \overset{\mathscr{M}g}{\nearrow} & \Downarrow \delta^{g,f} & \overset{\mathscr{M}f}{\searrow} & \\ \mathcal{H}(\mathbb{D})(C,C) & & \underset{\mathscr{M}(g\circ f)}{\searrow} & & \mathcal{H}(\mathbb{D})(A,A) \end{array}$$

for any $f : A \to B$ and $g : B \to C$, satisfying the commutativity of (2). Explicitely, each $\delta^{g,f}$ has components, for each horizontal 1-cell $J : C \nrightarrow C$,

$$\delta^{g,f}_J : (\mathscr{M}f \circ \mathscr{M}g)J \xrightarrow{\sim} \mathscr{M}(g\circ f)J = \breve{f} \odot \breve{g} \odot J \odot \widehat{g} \odot \widehat{f} \xrightarrow{\sim} \widetilde{gf} \odot J \odot \widehat{gf} \quad (24)$$

in $\mathcal{H}(\mathbb{D})(A,A)$, due to Lemma 3.7. Moreover, for any 0-cell A there is a natural isomorphism γ with components

$$\begin{array}{ccccc} & & \mathbf{1}_{\mathcal{H}(\mathbb{D})(A,A)} & & \\ & \overset{}{\nearrow} & \Downarrow \gamma^A & \overset{}{\searrow} & \\ \mathcal{H}(\mathbb{D})(A,A) & & \underset{\mathscr{M}(\mathrm{id}_A)}{\searrow} & & \mathcal{H}(\mathbb{D})(A,A) \end{array}$$

with components invertible arrows in $\mathcal{H}(\mathbb{D})(A,A)$

$$\gamma^X_G : G \xrightarrow{\sim} \mathscr{M}(\mathrm{id}_A)G = \widetilde{\mathrm{id}_A} \odot \widehat{\mathrm{id}_A} \quad (25)$$

again by Lemma 3.7; it can be verified that the axioms (3) are satisfied.

The Grothendieck category $\mathfrak{G}\mathscr{M}$ has as objects pairs (G, A) where A is a 0-cell and G is in $\mathcal{H}(\mathbb{D})(A,A)$, and as arrows $(\phi, f) : (G, A) \to (H, B)$ pairs

$$\begin{cases} G \xrightarrow{\phi} \check{f} \odot H \odot \hat{f} & \text{in } \mathcal{H}(\mathbb{D})(A,A) \\ A \xrightarrow{f} B & \text{in } \mathbb{D}_0. \end{cases}$$

It is not hard to verify its isomorphism with $\mathbb{D}^\bullet_1$: objects are the same (horizontal endo-1-cells), and there is a bijective correspondence between the morphisms, essentially given by (19). Given an arrow α_f in $\mathbb{D}^\bullet_1$, we obtain a composite 2-cell

$$\begin{array}{ccc} A & \overset{M}{\nrightarrow} & A \\ {\scriptstyle f}\downarrow & \Downarrow\alpha & \downarrow{\scriptstyle f} \\ B & \underset{N}{\nrightarrow} & B \end{array} \quad \mapsto \quad \begin{array}{ccccccc} A & \overset{1_A}{\nrightarrow} & A & \overset{M}{\nrightarrow} & A & \overset{1_A}{\nrightarrow} & A \\ {\scriptstyle \mathrm{id}_A}\downarrow & \Downarrow p_2 & \downarrow{\scriptstyle f} & \Downarrow\alpha & {\scriptstyle f}\downarrow & \Downarrow q_2 & \downarrow{\scriptstyle \mathrm{id}_A} \\ A & \underset{\hat{f}}{\nrightarrow} & B & \underset{N}{\nrightarrow} & B & \underset{\check{f}}{\nrightarrow} & B \end{array} \quad (26)$$

which is a morphism in $\mathfrak{G}\mathscr{M}$, where p_2 and q_2 come with the companion and conjoint as in Definition 3.4. This assignment is an isomorphism, with inverse mapping $\beta \mapsto (q_1 \odot 1_N \odot p_1)\beta$ for some $\beta : M \Rightarrow \check{f} \circ N \circ \hat{f}$ in $\mathcal{H}(\mathbb{D})(A,A)$. Thus $\mathscr{M}$ gives rise to a fibration $\mathfrak{G}\mathscr{M} \to \mathbb{D}_0$ which is isomorphic to $\mathbb{D}^\bullet_1 \to \mathbb{D}_0$ mapping G_X to X and α_f to f:

$$\begin{array}{ccc} \mathfrak{G}\mathscr{M} & \overset{\cong}{\longrightarrow} & \mathbb{D}^\bullet_1 \\ & \searrow \quad \swarrow & \\ & \mathbb{D}_0 & \end{array}$$

In a very similar way, it can be checked that $\mathscr{F}$ from (23) is a pseudofunctor, using again standard properties of companions and conjoints. Therefore, $\mathfrak{G}\mathscr{F} \cong \mathbb{D}_1^\bullet \to \mathbb{D}_0$ is an opfibration. □

Notice that $\mathbb{D}_1^\bullet$ being a bifibration over $\mathbb{D}_0$ could be deduced from the fibration part combined with Remark 2.20, since we have an adjunction $(\check{f} \odot \text{-} \odot \hat{f}) \vdash (\hat{f} \odot \text{-} \odot \check{f})$ for all f (Lemma 3.7).

Athough the above result was independently established as a generalization of our case study of (co)categories, as will be clear later, the fibration was also shown in [FGK11, Proposition 3.3], by restricting the cartesian liftings (19) of the fibration $(\mathfrak{s}, \mathfrak{t})\colon \mathbb{D}_1 \to \mathbb{D}_0$ to the category of endomorphisms, i.e.

$$\begin{aligned} \mathrm{Cart}(f, N) &= p_1 \odot 1_N \odot q_1 \colon \check{f} \odot N \odot \hat{f} \Rightarrow N \\ \mathrm{Cocart}(f, N) &= p_2 \odot 1_N \odot q_2 : N \Rightarrow \hat{f} \odot N \odot \check{f} \end{aligned} \tag{27}$$

Still, the pseudofunctor formulation of the above proof gives clearer perspectives for the objects involved in the applications.

We can now adjust the above constructions to obtain similar results for categories of monads and comonads in fibrant double categories. The following lemma ensures that is feasible.

Lemma 3.16. *For any vertical 1-cell $f\colon A \to B$ in a fibrant double category $\mathbb{D}$, the functors*

$$\begin{aligned} \check{f} \odot - \odot \hat{f} \colon \mathcal{H}(\mathbb{D})(B, B) &\longrightarrow \mathcal{H}(\mathbb{D})(A, A) \\ \hat{f} \odot - \odot \check{f} \colon \mathcal{H}(\mathbb{D})(A, A) &\longrightarrow \mathcal{H}(\mathbb{D})(B, B) \end{aligned}$$

are lax and colax monoidal respectively, for the monoidal endohom categories of the bicategory $\mathcal{H}(\mathbb{D})$ with horizontal composition.

Proof. For horizontal endo-1-cells $M, N : B \nrightarrow B$, the lax monoidal structure map

$$\phi_{M,N} : \check{f} \odot M \odot \hat{f} \odot \check{f} \odot N \odot \hat{f} \Rightarrow \check{f} \odot M \odot N \odot \hat{f}$$

is a natural transformation with components the composite 2-cells



where $\dot{\varepsilon}$ is the counit of the adjunction $\check{f} \dashv \hat{f}$. Similarly, ϕ_0 is given by



where $\dot{\eta}$ is the unit of $\check{f} \dashv \hat{f}$. These structure maps satisfy the usual conditions, hence $\check{f} \odot - \odot \hat{f}$ is a lax monoidal functor; dually, the colax monoidal structure of $\hat{f} \odot - \odot \check{f} \colon \mathcal{H}(\mathbb{D})(A, A) \to \mathcal{H}(\mathbb{D})(B, B)$ can be identified. □

Due to this lax and colax monoidal structure, the induced monoid structure of $(\check{f} \odot N \odot \hat{f})$ for a monoid $(\ N\colon B \nrightarrow B\ , \mu, \eta)$ and the induced comonoid structure of $(\hat{f} \odot C \odot \check{f})$ for $(\ C\colon A \nrightarrow A\ , \Delta, \epsilon)$ are



(28)



(29)

where $\dot{\eta}, \dot{\varepsilon}$ are the unit and counit of $\check{f} \dashv \hat{f}$.

Proposition 3.17. *If $\mathbb{D}$ is a fibrant double category,* $\mathbf{Mnd}(\mathbb{D})$ *is fibred over $\mathbb{D}_0$ and* $\mathbf{Cmd}(\mathbb{D})$ *is opfibred over $\mathbb{D}_0$.*

Proof. We will address the opfibration part; the fibration is established similarly. Once again, we will construct a pseudofunctor $\mathscr{K}\colon \mathbb{D}_0 \to \mathbf{Cat}$ for which the Grothendieck construction gives a total category isomorphic to $\mathbf{Cmd}(\mathbb{D})$, along with the evident forgetful functor to $\mathbb{D}_0$. This will resemble $\mathscr{F}$ from (23), but with fibre categories capturing the desired comonad structure.

An object A is mapped to the category $\mathbf{Comon}(\mathcal{H}(\mathbb{D})(A,A), \odot, 1_A)$ where double categorical monads with source and target A live (Remark 3.13). A vertical 1-cell $f\colon A \to B$ is mapped to the functor

$$\mathscr{K} f := \hat{f} \odot - \odot \check{f}\colon \mathbf{Comon}(\mathcal{H}(\mathbb{D})(A,A)) \to \mathbf{Comon}(\mathcal{H}(\mathbb{D})(B,B))$$

which is precisely the induced $\mathbf{Comon}(\mathscr{F} f)$ by Lemma 3.16. The fact that these data form a pseudofunctor follows in a straightforward way from $\mathscr{F}$ being a pseudofunctor; the natural isomorphisms δ and γ are given in a dual way to (24) and (25) using Lemma 3.7.

The induced Grothendieck category $\mathfrak{G}\mathscr{K}$ has as objects pairs (C, A) where $C \in \mathbf{Comon}(\mathcal{H}(\mathbb{D})(A,A))$ for a 0-cell A, and as arrows $(C, A) \to (D, B)$ pairs

$$\begin{cases} \hat{f} \odot C \odot \check{f} \xrightarrow{\psi} D & \text{in } \mathbf{Comon}(\mathcal{H}(\mathbb{D})(B,B)) \\ A \xrightarrow{f} B & \text{in } \mathbb{D}_0. \end{cases}$$

This category is isomorphic to $\mathbf{Cmd}(\mathbb{D})$ since they have the same objects, and there is a bijection between morphisms dually to (26)



where q_1, p_2 are as in Definition 3.4; checking that $p_1 \odot \alpha \odot q_1$ is a comonoid morphism follows from their properties. □

The above was again proved independently from the equivalent monad fibration proof of [FGK11, Proposition 3.3].

When $\mathbb{D}$ is moreover monoidal, these fibrations and opfibrations have a monoidal structure as well, in the sense of Definition 2.25.

Proposition 3.18. *Suppose that $\mathbb{D}$ is a fibrant monoidal double category. The bifibration $T\colon \mathbb{D}_1^\bullet \to \mathbb{D}_0$ as well as the fibration $S\colon \mathbf{Mnd}(\mathbb{D}) \to \mathbb{D}_0$ and opfibration $W\colon \mathbf{Cmd}(\mathbb{D}) \to \mathbb{D}_0$ are monoidal.*

Proof. By Proposition 3.14 and Definition 3.3 of a monoidal double category, all categories involved are monoidal. Moreover, for horizontal endo-1-cells $M\colon A \nrightarrow A$ and $N\colon B \nrightarrow B$,

$$T(M \otimes_1 N) = A \otimes_0 B = T(M) \otimes_1 T(N)$$

and similarly for P and W, by the mapping of $\otimes_1$ (18) whose restriction on $\mathbb{D}_1^\bullet, \mathbf{Mnd}(\mathbb{D}), \mathbf{Cmd}(\mathbb{D})$ is the tensor product.

Finally, $\otimes_1^\bullet\colon \mathbb{D}_1^\bullet \times \mathbb{D}_1^\bullet \to \mathbb{D}_1^\bullet$ preserves cartesian arrows: a pair of cartesian liftings $\mathrm{Cart}(f, M)$ and $\mathrm{Cart}(g, N)$ as in (27) is mapped to the top arrow


$(\hat{f} \odot M \odot \check{f}) \otimes_1 (\hat{g} \odot N \odot \check{g})$
$\mathrm{Cart}(f,M) \otimes_1 \mathrm{Cart}(g,N)$
$M \otimes_1 N$
$\cong$
$\mathrm{Cart}(f \otimes_0 g, M \otimes_1 N)$
$(\widehat{f \otimes_0 g}) \odot (M \otimes_1 N) \odot (\widetilde{f \otimes_0 g})$
in $\mathbb{D}_1^\bullet$
$A \otimes_0 C$
$B \otimes_0 C$
$f \otimes_0 g$
in $\mathbb{D}_0$

where the left side isomorphism is obtained by (17) and Lemma 3.7,

$$\Big((\hat{f} \odot M) \odot \check{f}\Big) \otimes_1 \Big((\hat{g} \odot N) \odot \check{g}\Big) \cong \Big((\hat{f} \odot M) \otimes_1 (\hat{g} \odot N)\Big) \odot (\check{f} \otimes_1 \check{g})$$
$$\cong (\hat{f} \otimes_1 \hat{g}) \odot (M \otimes_1 N) \odot (\check{f} \otimes_1 \check{g})$$

This vertical isomorphism can be shown to make the triangle commute hence the proof is complete, since this vertical isomorphism is reflected to monads and a dual cocartesian lifting triangle to comonads. □

3.3. **Locally closed monoidal double categories.** When $\mathbb{D}$ is a monoidal double category as in Definition 3.3, both vertical and horizontal categories are endowed with a monoidal structure, $(\mathbb{D}_0, \otimes_0, I)$ and $(\mathbb{D}_1, \otimes_1, 1_I)$. Naturally, one could expect that the appropriate notion of a *monoidal closed* double category would result in a similar 'local' closed structure for the two categories $\mathbb{D}_0$ and $\mathbb{D}_1$. For the usual definition though, this does not seem to be the case.

Definition 3.19. [GP04, §5] A *(weakly) monoidal closed* pseudo double category $\mathbb{D}$ is a monoidal double category such that each pseudo double functor $(\text{-} \otimes D) : \mathbb{D} \to \mathbb{D}$ has a lax right adjoint.

This definition uses *lax adjunctions* between pseudo double categories, as described in [GP04, 3.2]; double adjunctions are also studied in [FGK12]. Since these do not end up being relevant to the current work, we omit their details and simply discuss what they mean in this particular context.

The lax double functor $\text{-}\otimes D$ consists of the ordinary functors $(\text{-}\otimes_0 D\colon \mathbb{D}_0 \to \mathbb{D}_0, \text{-}\otimes_1 1_D\colon \mathbb{D}_1 \to \mathbb{D}_1)$. The existence of a lax right double adjoint, call it $\mathrm{Hom}^{\mathbb{D}}(D,-)$, amounts in particular to two ordinary adjunctions

$$\mathbb{D}_0 \underset{\mathrm{Hom}_0^{\mathbb{D}}(D,-)}{\overset{-\otimes_0 D}{\underset{\longleftarrow}{\overset{\longrightarrow}{\bot}}}} \mathbb{D}_0\,, \qquad \mathbb{D}_1 \underset{\mathrm{Hom}_1^{\mathbb{D}}(1_D,-)}{\overset{-\otimes_1 1_D}{\underset{\longleftarrow}{\overset{\longrightarrow}{\bot}}}} \mathbb{D}_1$$

for any 0-cell D in $\mathbb{D}$, such that conditions expressing compatibility with the horizontal composition and identities are satisfied. It immediately follows that $\mathbb{D}_0$ is a monoidal closed category; however this cannot be deduced for $\mathbb{D}_1$ as well, since 1_D is not an arbitrary horizontal 1-cell.

Due to the application of these notions to our context of interest later, we proceed to the definition of a different closed-like structure which arises naturally in what follows.

Definition 3.20. A monoidal (pseudo) double category $\mathbb{D}$ is called *locally monoidal closed* if it comes equipped with a lax double functor

$$H = (H_0, H_1)\colon \mathbb{D}^{\mathrm{op}} \times \mathbb{D} \longrightarrow \mathbb{D}$$

such that $\otimes_0 \dashv H_0$ and $\otimes_1 \dashv H_1$ are parametrized adjunctions.

By definition (16), the functor H_1 in particular is the mapping

$$H_1 :\ \mathbb{D}_1^{\mathrm{op}} \times \mathbb{D}_1 \longrightarrow \mathbb{D}_1 \tag{30}$$

$$\begin{array}{ccc}
(X \xrightarrow{M} Y, & Z \xrightarrow{N} W) & \longmapsto H_0(X,Z) \xrightarrow{H_1(M,N)} H_0(Y,W) \\
f\downarrow \quad \Downarrow\alpha \quad \downarrow g & h\downarrow \quad \Downarrow\beta \quad \downarrow k & H_0(f,h)\downarrow \quad \Downarrow H_1(\alpha,\beta) \quad \downarrow H_0(g,k) \\
(X' \xrightarrow[M']{} Y', & Z' \xrightarrow[N']{} W') & \longmapsto H_0(X',Z') \xrightarrow[H_1(M',N')]{} H_0(Y',W')
\end{array}$$

Call H the *internal hom* of $\mathbb{D}$. Clearly H_0 gives a monoidal closed structure on the vertical monoidal category $(\mathbb{D}_0, \otimes_0, I)$ and H_1 on the horizontal category $(\mathbb{D}_1, \otimes_1, 1_I)$. The above arguments justify that a monoidal closed structure on a double category does not imply a locally monoidal closed structure.

We will now explore the relation of this lax double functor H on a locally monoidal closed double category $\mathbb{D}$ with the categories of endomorphisms, monads and comonads discussed in Section 3.2. Recall that by Proposition 3.14, all these categories inherit a monoidal structure from $\mathbb{D}_1$.

Proposition 3.21. *Suppose $\mathbb{D}$ is a locally monoidal closed double category, with internal hom $H = (H_0, H_1)$. Then $H_1^\bullet$ endows the category of endomorphisms $\mathbb{D}_1^\bullet$ with a monoidal closed structure.*

Proof. The lax double functor H restricts to $H_1^\bullet\colon \mathbb{D}_1^{\bullet\mathrm{op}} \times \mathbb{D}_1^\bullet \to \mathbb{D}_1^\bullet$ by Corollary 3.11. The natural isomorphism $\mathbb{D}_1(M \otimes_1 N, P) \cong \mathbb{D}_1(M, H_1(N,P))$ which defines the adjunction $(-\otimes_1 N) \dashv H_1(N,-)$ for the monoidal closed category $\mathbb{D}_1$, considered only on endo-1-cells and endo-2-morphisms, implies that $\mathbb{D}_1^\bullet$ is also a monoidal closed category via $\otimes_1^\bullet \dashv H_1^\bullet$. □

The lax double functor $H\colon \mathbb{D}^{\mathrm{op}} \times \mathbb{D} \to \mathbb{D}$ by Proposition 3.12 induces

$$\mathbf{Mon}H : \mathbf{Cmd}(\mathbb{D})^{\mathrm{op}} \times \mathbf{Mnd}(\mathbb{D}) \to \mathbf{Mnd}(\mathbb{D}) \tag{31}$$

between the category of monads, since $\mathbf{Mnd}(\mathbb{D}^{\mathrm{op}}\times\mathbb{D}) \cong \mathbf{Mnd}(\mathbb{D}^{\mathrm{op}})\times\mathbf{Mnd}(\mathbb{D})$. This functor is fundamental in order to study enrichment relations between the category of monads and comonads in a symmetric locally monoidal closed (fibrant) double category, by applying results from Sections 2.3 and 2.4. For example, it is an action as explained below.

Lemma 3.22. *The functor* $\mathbf{Mon}H\colon \mathbf{Cmd}(\mathbb{D})^{\mathrm{op}} \times \mathbf{Mnd}(\mathbb{D}) \to \mathbf{Mnd}(\mathbb{D})$ *in a braided locally monoidal closed double category, as well as* $(\mathbf{Mon}H)^{\mathrm{op}}$*, is an action.*

Proof. The induced monoidal closed structure $H_1^\bullet$ on the braided monoidal $\mathbb{D}_1^\bullet$, as well as its opposite functor, are both actions of $\mathbb{D}_1^{\bullet\mathrm{op}}$ and $\mathbb{D}_1^\bullet$ respectively, as is the case in any braided monoidal closed category by Lemma 2.16. Since the forgetful functors from $\mathbf{Cmd}(\mathbb{D})$, $\mathbf{Mnd}(\mathbb{D})$ to $\mathbb{D}_1^\bullet$ reflect isomorphisms, the result follows. □

Theorem 3.23. *Suppose that* $(\mathbb{D}, \otimes, \mathbf{I})$ *is a symmetric locally monoidal closed double category, with internal hom* H*. If the induced functor* $\mathbf{Mon}H$ *has a parametrized right adjoint, then the category of monads* $\mathbf{Mnd}(\mathbb{D})$ *is enriched in the category of comonads* $\mathbf{Cmd}(\mathbb{D})$*.*

Moreover, if $\mathbf{Cmd}(\mathbb{D})$ *is monoidal closed this enrichment is cotensored, and if each* $\mathbf{Mon}H(M,-)$ *also has a right adjoint, the enrichment is tensored.*

Proof. The category of comonads is symmetric monoidal by Proposition 3.14. Since $\mathbf{Mon}H$ and $(\mathbf{Mon}H)^{\mathrm{op}}\colon \mathbf{Cmd}(\mathbb{D}) \times \mathbf{Mnd}(\mathbb{D})^{\mathrm{op}} \to \mathbf{Mnd}(\mathbb{D})^{\mathrm{op}}$ are actions, the existence of a right adjoint for the latter

$$S\colon \mathbf{Mnd}(\mathbb{D})^{\mathrm{op}} \times \mathbf{Mnd}(\mathbb{D}) \longrightarrow \mathbf{Cmd}(\mathbb{D})$$

induces the desired enrichment of $\mathbf{Mnd}(\mathbb{D})^{\mathrm{op}}$ and thus of $\mathbf{Mnd}(\mathbb{D})$, by Theorem 2.17. The rest of the clauses follow by assumption. □

Notice that monoidal closedness of $\mathbf{Cmd}(\mathbb{D})$ does not seem to follow in a straightforward way from that of $\mathbb{D}_1^\bullet$, Proposition 3.21. Even in the application that follows, this result is obtained after establishing (co)completeness and (co)monadicity properties for the specific structure, which are heavily related to the double category we choose to work in; see Proposition 4.33.

Moving to the fibrant case, we would now want to combine the above enrichment with the (op)fibration structure of monads and comonads over $\mathbb{D}_0$ to exhibit the formed picture as an enriched fibration, Definition 2.27. Working towards that end, notice that

$$\begin{array}{ccc} \mathbf{Cmd}(\mathbb{D}) \times \mathbf{Mnd}(\mathbb{D})^{\mathrm{op}} & \xrightarrow{(\mathbf{Mon}H)^{\mathrm{op}}} & \mathbf{Mnd}(\mathbb{D})^{\mathrm{op}} \\ \downarrow{\scriptstyle W\times S^{\mathrm{op}}} & & \downarrow{\scriptstyle S^{\mathrm{op}}} \\ \mathbb{D}_0 \times (\mathbb{D}_0)^{\mathrm{op}} & \xrightarrow[H_0^{\mathrm{op}}]{} & \mathbb{D}_0^{\mathrm{op}} \end{array} \tag{32}$$

commutes by definition of the involved functors, and moreover W, S^{op} are monoidal opfibrations by Proposition 3.18. Finally, the actions $(\mathbf{Mon}H)^{\mathrm{op}}$

and $(H_0)^{\mathrm{op}}$ are above each other in the sense of (13): by the mapping (30) that restricts as $\mathbf{Mon}H$ to (co)monads, the source and target of the action isomorphisms in $\mathbf{Mnd}(\mathbb{D})$ is precisely the action isomorphism in $\mathbb{D}_0$, e.g.

$$\begin{array}{ccc} H_0(X \otimes_0 Y, Z) & \xrightarrow{H_1(C \otimes_1 D, M)} & H_0(X \otimes_0 Y, Z) \\ \cong \downarrow & \Downarrow \cong & \downarrow \cong \\ H_0(X, H_0(Y, Z)) & \xrightarrow[H_1(C, H_1(D, M))]{} & H_0(X, H_0(Y, Z)) \end{array}$$

As a result, when $\mathbf{Mon}H$ preserves cartesian liftings, then W coacts on S^{op} in the sense of Definition 2.26. We can now apply the dual of Theorem 2.29.

Theorem 3.24. *Suppose $\mathbb{D}$ is a symmetric locally monoidal closed fibrant double category. If $(\mathbf{Mon}H)^{\mathrm{op}}$ preserves cocartesian liftings, and (32) has a general lax parametrized opfibred adjoint as in Definition 2.24, then the fibration $S\colon \mathbf{Mnd}(\mathbb{D}) \to \mathbb{D}_0$ is enriched in the symmetric monoidal opfibration $W\colon \mathbf{Cmd}(\mathbb{D}) \to \mathbb{D}_0$.*

4. Enriched matrices, categories and cocategories

In this section, we initially study the double category of $\mathcal{V}$-matrices, also briefly considered in [Shu08]. After establishing its monoidal fibrant double categorical structure, special focus will be given to its well-known horizontal bicategory of enriched matrices. The main references for that are [BCSW83] and [KL01]; in the former, the more general bicategory $\mathcal{W}$-**Mat** of matrices enriched in a bicategory $\mathcal{W}$ was studied, leading to the theory of bicategory enriched categories. For the one-object case, *i.e.* monoidal categories, see [Bén73] and [Wol74].

Furthermore, we investigate the categories of monads and comonads in the double category $\mathcal{V}$-$\mathbb{M}$**at**. These are specifically $\mathcal{V}$-**Cat** of $\mathcal{V}$-enriched categories and functors [Kel05], and $\mathcal{V}$-**Cocat** of $\mathcal{V}$-enriched *cocategories* and *cofunctors*. These categories are studied in detail, passing from the double categorical to the bicategorical view of $\mathcal{V}$-matrices according to our needs. Finally, employing previous results regarding enrichment relations of these categories, the goal is to establish an enrichment of $\mathcal{V}$-categories in $\mathcal{V}$-cocategories.

As an intermediate step, $\mathcal{V}$-enriched graphs are given special attention; they are the category of endomorphisms and they provide a natural framework for categories and cocategories, which are enriched graphs with extra structure. As a primary reference, in [ML98, 11.7] the notion of a small (directed) graph was employed to describe the free category construction, in analogy with the free monoid construction on a set. Moreover, the idea of O-graphs with a fixed set of objects O inspires the fibrational view of these categories. For the main results regarding $\mathcal{V}$-**Grph** and $\mathcal{V}$-**Cat** from a more traditional point of view, Wolff's [Wol74] is a classical reference for the case of a symmetric monoidal closed category, whereas for the description of $\mathcal{V}$-graphs in terms of $\mathcal{V}$-matrices, we again closely follow [BCSW83, KL01].

There is a 2-dimensional aspect for all the basic categories we study in this chapter, e.g. $\mathcal{V}$-natural transformations. However, we choose to omit its description in this treatment, because it is not necessary for the current level of our main objectives.

4.1. **$\mathcal{V}$-matrices.** Suppose that $\mathcal{V}$ is a monoidal category with coproducts which are preserved by the tensor product functor $- \otimes -$ in each variable; for example, this is certainly the case when $\mathcal{V}$ is monoidal closed.

For sets X and Y, a *$\mathcal{V}$-matrix* $S : X \nrightarrow Y$ from X to Y is defined to be a functor $S : X \times Y \to \mathcal{V}$, where the set $X \times Y$ is viewed as a discrete category. This can equivalently given by a family of objects in $\mathcal{V}$

$$\{S(x, y)\}_{(x,y)\in X\times Y}$$

sometimes also denoted as $\{S_{x,y}\}_{X\times Y}$. For example, each set X gives rise to a $\mathcal{V}$-matrix $1_X : X \nrightarrow X$ called the *identity matrix* given by

$$1_X(x, x') = \begin{cases} I, & \text{if } x = x' \\ 0, & \text{otherwise} \end{cases}$$

where I is the unit object in $\mathcal{V}$ and 0 is the initial object.

There is a double category $\mathcal{V}$-$\mathbb{M}\mathbf{at}$ of $\mathcal{V}$-matrices as in Definition 3.1, with vertical category $\mathcal{V}$-$\mathbb{M}\mathbf{at}_0$ the usual category of sets and functions **Set**. Its horizontal category $\mathcal{V}$-$\mathbb{M}\mathbf{at}_1$ consists of $\mathcal{V}$-matrices $S : X \nrightarrow Y$ as horizontal 1-cells, and 2-morphisms ${}^f\alpha^g : S \Rightarrow T$ are natural transformations



(33)

given by families of arrows $\alpha_{x,y} : S(x, y) \to T(fx, gy)$ in $\mathcal{V}$, for all $x \in X$ and $y \in Y$. There is a functor $\mathbf{1} \colon \mathcal{V}\text{-}\mathbb{M}\mathbf{at}_0 \to \mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1$ which gives the identity $\mathcal{V}$-matrix $1_X : X \nrightarrow X$ for each X, and the unit 2-morphism 1_f with components

$$(1_f)_{x,x'} : 1_X(x, x') \to 1_X(x, x') \equiv \begin{cases} I \xrightarrow{1_I} I, & \text{if } x = x' \\ 0 \to 0, & \text{if } x \neq x'. \end{cases}$$

The source and target functors give the evident sets and functions, and the horizontal composition functor

$$\odot : \mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1 \times_{\mathcal{V}\text{-}\mathbb{M}\mathbf{at}_0} \mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1 \to \mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1$$

maps two composable $\mathcal{V}$-matrices $T : Y \nrightarrow Z$ and $S : X \nrightarrow Y$ to the matrix $T \circ S : X \nrightarrow Z$, given by the family of objects in $\mathcal{V}$

$$(T \circ S)(x, z) = \sum_{y\in Y} S(x, y) \otimes T(y, z) \tag{34}$$

for all $z \in Z$ and $x \in X$, reminiscent of the usual matrix multiplication. The horizontal composite of 2-morphisms ${}^f(\beta \odot \alpha)^g : T \circ S \Rightarrow T' \circ S'$ as in (15) is given by the composite arrows

$$\begin{array}{ccc} \sum_y S(x, y) \otimes T(y, z) & \xrightarrow{\sum \alpha_{x,y}\otimes\beta_{y,z}} & \sum_y S'(fx, gy) \otimes T'(gy, hz) \\ & \searrow & \downarrow \\ & & \sum_{y'} S'(fx, y') \otimes T'(y', hz) \end{array} \tag{35}$$

in $\mathcal{V}$, for all $x \in X$ and $z \in Z$. Compatibility conditions of source and target functors with composition can be easily checked.

For composable $\mathcal{V}$-matrices $X \overset{S}{\nrightarrow} Y \overset{T}{\nrightarrow} Z \overset{R}{\nrightarrow} W$, the associator α has components globular isomorphisms $\alpha^{R,T,S} : (R \odot T) \odot S \xrightarrow{\sim} R \odot (T \odot S)$ given by the family $\{\alpha_{x,w}\}$ of composite isomorphisms

$$\begin{array}{ccc} \sum_z \left(\sum_y S_{x,y} \otimes T_{y,z} \right) \otimes R_{z,w} & \dashrightarrow & \sum_y S_{x,y} \otimes \left(\sum_z T_{y,z} \otimes R_{z,w} \right) \\ \downarrow \cong & & \uparrow \cong \\ \sum_{y,z} \left((S_{x,y} \otimes T_{y,z}) \otimes R_{z,w} \right) & \xrightarrow[\sum a]{} & \sum_{y,z} \left(S_{x,y} \otimes (T_{y,z} \otimes R_{z,w}) \right) \end{array}$$

where a is the associativity constraint of $\mathcal{V}$ and the invertible arrows express the fact that $\otimes$ commutes with colimits. Finally, for each $\mathcal{V}$-matrix $S : X \nrightarrow Y$, the unitors λ, ρ have components globular $\lambda^S : 1_Y \odot S \xrightarrow{\sim} S, \rho^S : S \odot 1_X \xrightarrow{\sim} S$ given by families

$$\lambda^S_{x,y} : \sum_{y' \in Y} S(x,y') \otimes 1_Y(y',y) \equiv S(x,y) \otimes I \xrightarrow{r_{S(x,y)}} S(x,y)$$

$$\rho^S_{x,y} : \sum_{x' \in X} 1_X(x,x') \otimes S(x',y) \equiv I \otimes S(x,y) \xrightarrow{l_{S(x,y)}} S(x,y).$$

of morphisms in $\mathcal{V}$. The respective coherence conditions are satisfied, thus these data indeed define a double category.

Moreover, $\mathcal{V}$-$\mathbb{M}\mathbf{at}$ is fibrant as in Definition 3.5. Any function $f : X \to Y$ determines two $\mathcal{V}$-matrices, $f_* : X \nrightarrow Y$ and $f^* : Y \nrightarrow X$, given by

$$f_*(x,y) = f^*(y,x) = \begin{cases} I, & \text{if } f(x) = y \\ 0, & \text{otherwise} \end{cases} \tag{36}$$

These are precisely the companion and the conjoint of the vertical 1-cell f, since they come equipped with appropriate 2-cells as in Definition 3.4:

$$\begin{array}{ccc} X & \overset{f_*}{\nrightarrow} & Y \\ {\scriptstyle f}\downarrow & \Downarrow p_1 & \| \\ Y & \underset{1_Y}{\nrightarrow} & Y \end{array} \qquad \begin{array}{ccc} X & \overset{1_X}{\nrightarrow} & X \\ \| & \Downarrow p_2 & \downarrow{\scriptstyle f} \\ X & \underset{f_*}{\nrightarrow} & Y \end{array} \qquad \text{are given by}$$

$$f_*(x,y) \xrightarrow{(p_1)_{x,y}} 1_Y(fx,y) = \begin{cases} I \xrightarrow{\mathrm{id}} I, & \text{if } y = fx \\ 0 \xrightarrow{\mathrm{id}} 0, & \text{otherwise} \end{cases}$$

$$1_X(x,x') \xrightarrow{(p_2)_{x,x'}} f_*(x,fx') = \begin{cases} I \xrightarrow{\mathrm{id}} I, & \text{if } x = x' \\ 0 \xrightarrow{!} \begin{cases} I, & fx = fx' \\ 0, & \text{else} \end{cases} & \text{if } x \neq x' \end{cases}$$

satisfying the required relations, and similarly for f^*.

When $\mathcal{V}$ is braided monoidal, the double category $\mathcal{V}$-$\mathbb{M}\mathbf{at}$ has a monoidal structure as in Definition 3.3. The required double functors $\otimes = (\otimes_0, \otimes_1)$ and

$\mathbf{I} = (\mathbf{I}_0, \mathbf{I}_1)$ consist of the cartesian monoidal structure on the vertical category $(\mathbf{Set}, \times, \{*\})$ and

$$\otimes_1 : \ \mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1 \times \mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1 \longrightarrow \mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1 \tag{37}$$

$$\begin{array}{ccccc} (X & \xrightarrow{S} & Y & , & Z \xrightarrow{T} W) \\ {\scriptstyle f}\downarrow & \Downarrow\alpha & \downarrow{\scriptstyle g} & & {\scriptstyle h}\downarrow \ \Downarrow\beta \ \downarrow{\scriptstyle k} \\ (X' & \xrightarrow[S']{} & Y' & , & Z' \xrightarrow[T']{} W') \end{array} \mapsto \begin{array}{ccc} X \times Z & \xrightarrow{S\otimes T} & Y \times W \\ {\scriptstyle f\times h}\downarrow & \Downarrow\alpha\otimes\beta & \downarrow{\scriptstyle g\times k} \\ X' \times Z' & \xrightarrow[S'\otimes T']{} & Y' \times W' \end{array}$$

which is defined on objects and morphisms by the families in $\mathcal{V}$

$$(S \otimes T)\left((x,z),(y,w)\right) := S(x,y) \otimes T(z,w) \tag{38}$$

$$(\alpha \otimes \beta)_{(x,z),(y,w)} := S(x,y) \otimes T(z,w) \xrightarrow{\alpha_{x,y}\otimes\beta_{z,w}} S'(fx,gy) \otimes T'(hz,kw).$$

Along with the $\mathcal{V}$-matrix $\mathcal{I} : \{*\} \nrightarrow \{*\}$ given by $\mathcal{I}(*,*) = I_{\mathcal{V}}$, this defines a monoidal structure of $\mathcal{V}$-$\mathbb{M}\mathbf{at}_1$. The conditions for $\mathfrak{s}$ and $\mathfrak{t}$ are satisfied, and the globular isomorphisms (17) come down to the tensor product in $\mathcal{V}$ commuting with coproducts. More specifically, for matrices $S\colon X \nrightarrow Y$, $T\colon Z \nrightarrow W$, $S'\colon Y \nrightarrow U$, $T'\colon W \nrightarrow V$ we can compute the isomorphic families in $\mathcal{V}$

$$\left((S \otimes_1 T) \odot (S' \otimes_1 T')\right)_{(x,z),(u,v)} = \sum_{(y,w)} S_{x,y} \otimes T_{z,w} \otimes S'_{y,u} \otimes T'_{w,v}$$

$$\left((S \odot S') \otimes_1 (T \odot T')\right)_{(x,z),(u,v)} = \sum_{y} \left(S_{x,y} \otimes S'_{y,u}\right) \otimes \sum_{w} \left(T_{z,w} \otimes T'_{w,v}\right)$$

via the braiding, and also $1_{X\times Y} \cong 1_X \otimes_1 1_Y$ in a straightforward way. This monoidal structure on $\mathcal{V}$-$\mathbb{M}\mathbf{at}$ is symmetric, when the base monoidal category $\mathcal{V}$ is symmetric. Then $\mathbf{Set}$ and $\mathcal{V}$-$\mathbb{M}\mathbf{at}_1$ are both symmetric monoidal categories, and the rest of the axioms follow.

Proposition 4.1. *If $\mathcal{V}$ is a monoidal category with coproducts, such that the tensor product preserves them in both variables, the double category $\mathcal{V}$-$\mathbb{M}\mathbf{at}$ is fibrant. Moreover, if $\mathcal{V}$ is braided monoidal then $\mathcal{V}$-$\mathbb{M}\mathbf{at}$ is monoidal, and if $\mathcal{V}$ is symmetric then so is $\mathcal{V}$-$\mathbb{M}\mathbf{at}$.*

Remark 4.2. It is evident that there is a strong relation between $\mathcal{V}$-$\mathbb{M}\mathbf{at}$ and $\mathcal{V}$-$\mathbb{P}\mathbf{rof}$, the double category of $\mathcal{V}$-profunctors. On a first level, we can see that $\mathcal{V}$-matrices are special cases of $\mathcal{V}$-profunctors when the latter are considered only on discrete categories. In [Shu08, 11.8], a more elaborate relation between these double categories is established: $\mathbb{M}\mathbf{od}(\mathcal{V}\text{-}\mathbb{M}\mathbf{at}) = \mathcal{V}\text{-}\mathbb{P}\mathbf{rof}$ for a construction $\mathbb{M}\mathbf{od}$ building new fibrant double categories from old, with vertical category that of monads.

When $\mathcal{V}$ is moreover monoidal closed with products, we can determine a locally monoidal closed structure on $\mathcal{V}$-$\mathbb{M}\mathbf{at}$. Following Definition 3.20, we are after a lax double functor

$$H = (H_0, H_1)\colon \mathcal{V}\text{-}\mathbb{M}\mathbf{at}^{\mathrm{op}} \times \mathcal{V}\text{-}\mathbb{M}\mathbf{at} \to \mathcal{V}\text{-}\mathbb{M}\mathbf{at} \tag{39}$$

which endows $\mathcal{V}\text{-}\mathbb{M}\mathbf{at}_0 = \mathbf{Set}$ and $\mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1$ with a monoidal closed structure. For the vertical category, we clearly have the exponentiation functor

$$H_0 : \mathbf{Set}^{\mathrm{op}} \times \mathbf{Set} \xrightarrow{(-)^{(-)}} \mathbf{Set}$$

as the internal hom. For the horizontal category, we can define

$$H_1 : \ \mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1^{\mathrm{op}} \times \mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1 \longrightarrow \mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1 \tag{40}$$

$$\begin{array}{ccccc}
(X & \xrightarrow{S} & Y & , & Z \xrightarrow{T} W) \mapsto Z^X \xrightarrow{H_1(S,T)} W^Y \\
{\scriptstyle f}\downarrow & \Downarrow\alpha & \downarrow{\scriptstyle g} & & {\scriptstyle h}\downarrow \ \Downarrow\beta \ \downarrow{\scriptstyle k} \qquad {\scriptstyle h^f}\downarrow \ \Downarrow H_1(\alpha,\beta) \downarrow{\scriptstyle k^g} \\
(X' & \xrightarrow[S']{} & Y' & , & Z' \xrightarrow[T']{} W') \mapsto Z'^{X'} \xrightarrow[H_1(S',T')]{} W'^{Y'}
\end{array}$$

on horizontal 1-cells given by families of objects in $\mathcal{V}$, for $n \in Z^X, m \in W^Y$,

$$H_1(S,T)(n,m) = \prod_{x,y} [S(x,y), T(n(x), m(y))] \tag{41}$$

and on 2-morphisms $H_1(\alpha,\beta) : H_1(S,T)(n,m) \to H_1(S',T')(h^f(n), k^g(m))$ by families of arrows in $\mathcal{V}$

$$\prod_{x,y} [S(x,y), T(n(x), m(y))] \to \prod_{x',y'} [S'(x',y'), T'(hnf(x'), kmg(y'))] \tag{42}$$

which correspond under $(\text{-} \otimes X) \dashv [X, -]$ in $\mathcal{V}$ for fixed x', y' to the composite

$$\begin{array}{ccc}
\prod_{x,y}[S(x,y), T(nx,my)] \otimes S'(x',y') & \dashrightarrow & T'(hnfx', kmgy') \\
{\scriptstyle 1\otimes\alpha_{x',y'}}\downarrow & & \uparrow{\scriptstyle \beta_{nfx',mgy'}} \\
\prod_{x,y}[S(x,y), T(nx,my)] \otimes S(fx',gy') & & \\
{\scriptstyle \pi_{fx',gy'}\otimes 1}\downarrow & & \\
[S(fx',gy'), T(nfx',mgy')] \otimes S(fx',gy') & \xrightarrow{\mathrm{ev}} & T(nfx',mgy').
\end{array}$$

The globular transformations from Definition 3.2 for a lax double functor

$$\begin{array}{ccccc}
 & {\scriptstyle H_1(S,T)} & W^Z & {\scriptstyle H_1(R,O)} & \\
Y^X & & \Downarrow \delta_{(S,T),(R,O)} & & V^U \\
 & & {\scriptstyle H_1(R\circ S, O\circ T)} & &
\end{array}
\qquad\qquad
\begin{array}{ccc}
 & {\scriptstyle 1_{Y^X}} & \\
Y^X & \Downarrow \gamma_{(X,Y)} & Y^X \\
 & {\scriptstyle H_1(1_X, 1_Y)} &
\end{array}$$

for each $(R : Z \nrightarrow U,\ O : W \nrightarrow V)$ and $(S : X \nrightarrow Z,\ T : Y \nrightarrow W)$ are given by families of arrows in $\mathcal{V}$

$$\sum_{q \in W^Z} H_1(S,T)(k,q) \otimes H_1(R,O)(t,q) \xrightarrow{\delta_{k,t}} \prod_{(x,u)} [(R\circ S)(x,u), (O\circ T)(kx,tu)]$$

$$\gamma_{k,k} : I \xrightarrow{\sim} [1_X(x,x), 1_Y(kx,kx)] = [I,I]$$

for all $k \in Y^X, t \in V^U$, $x = x' \in X$. These again can be understood via their transposes under the tensor-hom adjunction, *i.e.* composites of projections, inclusions, braidings and evaluations, using the fact that the tensor product

preserves sums. The coherence axioms of Definition 2.2 as for a lax functor of bicategories are satisfied, therefore $H = (H_0, H_1)$ is a lax double functor.

Proposition 4.3. *Under the above assumptions, the functor H_1 (40) constitutes a monoidal closed structure for $(\mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1, \otimes_1, 1_I)$.*

Proof. We need to show that $- \otimes_1 T \dashv H_1(T, -) \colon \mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1 \to \mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1$ for any $\mathcal{V}$-matrix $T \colon Z \nrightarrow W$, i.e. there is a natural bijection between 2-morphisms

$$\begin{array}{ccc} X \times Z & \xrightarrow{S \otimes_1 T} & Y \times W \\ {\scriptstyle f}\downarrow & \Downarrow \alpha & \downarrow{\scriptstyle g} \\ U & \xrightarrow[P]{} & V \end{array} \quad \text{and} \quad \begin{array}{ccc} X & \xrightarrow{S} & Y \\ {\scriptstyle k}\downarrow & \Downarrow \beta & \downarrow{\scriptstyle l} \\ U^Z & \xrightarrow[H_1(T,P)]{} & V^W \end{array}$$

Taking components of the left side 2-morphism (33) and using the monoidal closed structure of $\mathcal{V}$, we deduce that any $\alpha_{(x,z),(y,x)} \colon S(x,y) \otimes T(z,w) \to P(f(x,z), g(y,w))$ bijectively corresponds, for all $x \in X, y \in Y, z \in Z, w \in W$, to some $\mathcal{V}$-morphism $S(x,y) \to [T(z,w), P(f(x,z), g(y,w))]$. Since $\mathcal{V}$ has products and **Set** is closed, this uniquely corresponds to some

$$\beta_{x,y} \colon S(x,y) \to \prod_{z,w} [T(z,w), P(f_x(z), g_y(w))]$$

for the transpose functions $f_x \in U^X$, $g_y \in V^W$ and the proof is complete. □

Hence we built a lax double functor H which satisfies the conditions of Definition 3.20; the desired result follows.

Corollary 4.4. *If $\mathcal{V}$ is a braided monoidal closed category with products and coproducts, the monoidal double category $\mathcal{V}$-$\mathbb{M}\mathbf{at}$ is locally monoidal closed.*

The horizontal bicategory $\mathcal{H}(\mathcal{V}\text{-}\mathbb{M}\mathbf{at})$ of the double category of $\mathcal{V}$-matrices is the well-known bicategory $\mathcal{V}$-**Mat**. In more detail, sets and $\mathcal{V}$-matrices are the 0- and 1-cells, and 2-cells between $\mathcal{V}$-matrices S and S' are globular 2-morphisms, i.e. natural transformations

$$X \;\begin{array}{c} \xrightarrow{S} \\ \Downarrow \sigma \\ \xrightarrow[S']{} \end{array}\; Y \;:=\; X \times Y \;\begin{array}{c} \xrightarrow{S} \\ \Downarrow \sigma \\ \xrightarrow[S']{} \end{array}\; \mathcal{V}$$

given by families $\sigma_{x,y} : S(x,y) \to S'(x,y)$ of arrows in $\mathcal{V}$. The horizontal composition $\circ : \mathcal{V}\text{-}\mathbf{Mat}(Y,Z) \times \mathcal{V}\text{-}\mathbf{Mat}(X,Y) \to \mathcal{V}\text{-}\mathbf{Mat}(X,Z)$ is given by matrix multiplication (34) on objects, and by the special case of (35) mapping (σ, τ) to $\{(\tau * \sigma)_{x,z}\} = \{\sum \sigma_{x,y} \otimes \tau_{y,z}\}$ on morphisms.

Many useful properties of the companion and conjoint (36) can be deduced in the bicategorical context from Lemma 3.7. For example, for any $f \colon X \to Y$ we have an adjunction $f_* \dashv f^*$ in the bicategory $\mathcal{V}$-**Mat**, with unit and counit

$$X \;\begin{array}{c} \xrightarrow{1_X} \\ \Downarrow \dot{\eta} \\ \xrightarrow[f^* \circ f_*]{} \end{array}\; X \qquad \text{and} \qquad Y \;\begin{array}{c} \xrightarrow{f_* \circ f^*} \\ \Downarrow \dot{\varepsilon} \\ \xrightarrow[1_Y]{} \end{array}\; Y \tag{43}$$

with components arrows in $\mathcal{V}$

$$\dot{\varepsilon}_{y,y'} : (f_* \circ f^*)(y,y') \to 1_Y(y,y') \equiv \begin{cases} \sum\limits_{x \in f^{-1}(y)} I \otimes I \xrightarrow{r_I} I, & \text{if } y = y' \\ 0 \xrightarrow{!} 0, & \text{if } y \neq y' \end{cases}$$

$$\dot{\eta}_{x,x'} : 1_X(x,x') \to (f^* \circ f_*)(x',x) \equiv \begin{cases} I \xrightarrow{(r_I)^{-1}} I \otimes I, & \text{if } x' = x \\ 0 \xrightarrow{!} \begin{cases} I \otimes I, & fx = fx' \\ 0, & \text{else} \end{cases} & \text{if } x' \neq x \end{cases}$$

Notice that $\dot{\eta}$ and $\dot{\varepsilon}$ are isomorphisms if and only if f is a bijection.

For explicit calculations in the context of $\mathcal{V}$-matrices, it will be useful to compute that for any $F\colon X \to Y$, $S : Y \nrightarrow Z$ and $T : Z \nrightarrow Y$,

$$(S \circ f_*)(x,z) = \sum_{y \in Y} f_*(x,y) \otimes S(y,z) = I \otimes S(fx,z) \overset{r}{\cong} S(fx,z) \tag{44}$$

$$(f^* \circ T)(z,x) = \sum_{y \in Y} T(z,y) \otimes f^*(y,x) = T(z,fx) \otimes I \overset{l}{\cong} T(z,fx)$$

are the families in $\mathcal{V}$ that define the composite matrices $S \circ f_*$ and $f^* \circ T$. Using such machinery, we re-obtain the following results for composites of companions and conjoints, Item 2.

Lemma 4.5. *Let $f : X \to Y$ and $g : Y \to Z$ be functions. There exist isomorphisms*

$$\zeta^{g,f} : g_* \circ f_* \cong (gf)_* : X \longrightarrow\!\!\!\!\!\rightarrow Z$$
$$\xi^{g,f} : f^* \circ g^* \cong (gf)^* : Z \longrightarrow\!\!\!\!\!\rightarrow X$$

which are families of invertible arrows

$$\zeta^{g,f}_{x,z} = \xi^{g,f}_{z,x} : \begin{cases} I \otimes I \xrightarrow{r_I = l_I} I, & \text{if } g(f(x)) = z \\ 0 \xrightarrow{!} 0, & \text{otherwise} \end{cases} \tag{45}$$

Under the assumptions of Proposition 4.1, $\mathcal{V}$-$\mathbb{M}$**at** is a fibrant monoidal double category therefore Theorem 3.8 applies.

Proposition 4.6. *If $\mathcal{V}$ is a braided monoidal category with coproducts such that $\otimes$ preserves them in both enries, the bicategory $\mathcal{V}$-**Mat** is a monoidal bicategory; if $\mathcal{V}$ is symmetric then so is $\mathcal{V}$-**Mat**.*

The monoidal unit is the unit $\mathcal{V}$-matrix $\mathcal{I}$ and the induced tensor product pseudofunctor $\otimes : \mathcal{V}\text{-}\mathbf{Mat} \times \mathcal{V}\text{-}\mathbf{Mat} \to \mathcal{V}\text{-}\mathbf{Mat}$ maps two sets X, Y to their cartesian product $X \times Y$, and the functor

$$\otimes_{(X,Y),(Z,W)} : \mathcal{V}\text{-}\mathbf{Mat}(X,Z) \times \mathcal{V}\text{-}\mathbf{Mat}(Y,W) \to \mathcal{V}\text{-}\mathbf{Mat}(X \times Y, Z \times W),$$

is defined as in (37) for globular 2-morphisms.

When $\mathcal{V}$ is moreover monoidal closed with products, its locally monoidal structure $H = (H_0, H_1)$ (39) induces a lax functor of bicategories

$$\mathrm{Hom} : (\mathcal{V}\text{-}\mathbf{Mat})^{\mathrm{co}} \times \mathcal{V}\text{-}\mathbf{Mat} \longrightarrow \mathcal{V}\text{-}\mathbf{Mat}$$

where $\mathcal{V}$-$\mathbf{Mat}^{\mathrm{co}}$ is the bicategory of $\mathcal{V}$-matrices with reversed 2-cells, since $\mathcal{H}(\mathbb{D}^{\mathrm{op}}) = (\mathcal{H}(\mathbb{D}))^{\mathrm{co}}$. On objects is given by exponentiation, and the functor on hom-categories, for all $(X,Y),(Z,W)$, is

$$\mathrm{Hom}_{(X,Y),(Z,W)} \colon \mathcal{V}\text{-}\mathbf{Mat}(X,Z)^{\mathrm{op}} \times \mathcal{V}\text{-}\mathbf{Mat}(Y,W) \to \mathcal{V}\text{-}\mathbf{Mat}(Y^X, W^Z) \quad (46)$$

is given by $\mathrm{Hom}(S,T) = H_1(S,T)$ as in (41) on objects and $\mathrm{Hom}(\sigma,\tau) = H_1(\sigma,\tau)$ as in (42) on globular 2-morphisms, i.e. 2-cells.

The hom-categories of the bicategory $\mathcal{V}$-$\mathbf{Mat}$ are the functor categories $\mathcal{V}$-$\mathbf{Mat}(X,Y) = \mathcal{V}^{X\times Y}$. The endo-hom-categories for a fixed set X will play an important role; the following proposition exhibits some useful properties.

Proposition 4.7. *Let $\mathcal{V}$ be a monoidal category with all colimits such that $\otimes$ preserves them on both entries. The category $\mathcal{V}$-$\mathbf{Mat}(X,X) = [X\times X, \mathcal{V}]$ for any set X*

(i) is cocomplete and has all limits that exist in $\mathcal{V}$;
(ii) is a monoidal category, and $\otimes = \circ$ preserves any colimit on both entries;
(iii) is locally presentable when $\mathcal{V}$ is;
(iv) is monoidal closed when $\mathcal{V}$ is monoidal closed with products.

Proof.
(i) They are formed pointwise from those in $\mathcal{V}$.
(ii) $(\mathcal{V}\text{-}\mathbf{Mat}(X,X), \circ, 1_X)$ is monoidal for any bicategory, Remark 2.9.
If $(G_j \to G \,|\, j \in \mathcal{J})$ is a colimiting cocone of shape $\mathcal{J}$ in $\mathcal{V}$-$\mathbf{Mat}(X,X)$, for any $x,y \in X$ the arrows $G_j(x,y) \to G(x,y)$ form colimiting cocones in $\mathcal{V}$. If we apply $S \circ -$, we obtain a collection of 2-cells $(S\circ G_j \to S \circ G \,|\, j \in \mathcal{J})$ in $\mathcal{V}$-$\mathbf{Mat}$. For this to be a colimit, for any $x,z \in X$ the arrows

$$\sum_{y\in X} G_j(x,y) \otimes S(y,z) \longrightarrow \sum_{y\in X} \mathrm{colim}_j G_j(x,y) \otimes S(y,z)$$

must be colimiting in $\mathcal{V}$, which is the case since $(-\otimes A)$ preserves colimits:

$$\begin{aligned}\sum_{y\in X} (\mathrm{colim}_j G_j(x,y)) \otimes S(y,z) &\cong \sum_{y\in X} \mathrm{colim}_j (G_j(x,y) \otimes S(y,z)) \\ &\cong \mathrm{colim}_j (\sum_{y\in X} G_j(x,y) \otimes S(y,z)).\end{aligned}$$

(iii) This follows from [AR94, 1.54]: any functor category $[\mathcal{A},\mathcal{V}]$ over a presentable category $\mathcal{V}$ is also presentable.
(iv) This is obtained by a restriction of (41) on globular 2-morphisms. It is not hard to establish a bijective correspondence

$$\frac{S\circ T \longrightarrow R \qquad \text{in } \mathcal{V}\text{-}\mathbf{Mat}(X,X)}{S \longrightarrow F(T,R) \qquad \text{in } \mathcal{V}\text{-}\mathbf{Mat}(X,X)}$$

for $G(T,R)(x,y) := \prod_{z\in X} [T(y,z), R(x,z)]$. □

Remark 4.8. The endo-hom-categories of $\mathcal{V}$-$\mathbf{Mat}$ have in fact a *duoidal* structure, since they are equipped with a second, pointwise monoidal product

$$(S \bullet T)(x,y) := S(x,y) \otimes T(x,y), \quad J(x,y) = I,$$

This point of view is discussed in [BCV16, §7] without explicitly making use of $\mathcal{V}$-matrices, and is crucial concerning the expression of the so-called *semi-Hopf categories*, i.e. categories enriched in coalgebras, as bimonoids in $(\mathcal{V}\text{-}\mathbf{Mat}(X,X), \circ, \bullet, 1_X, J)$ and subsequently of *Hopf categories* as Hopf monoids in that duoidal category. This approach is relevant to work in progress regarding the expression of Hopf categories as *Hopf monads* in a bicategorical or even a double categorical context. Work in similar direction, establishing an abstract context for various Hopf-structure generalized notions, are [BL16, B16]; in fact, the above duoidal structure can be seen as a consequence of all sets being *map comonoidales* in the monoidal bicategory $\mathcal{V}$-**Mat**.

In Section 2.2, the categories of monoids and comonoids of locally presentable monoidal categories were considered. Due to the first three part of the above proposition, we obtain the following corollary to Proposition 2.13.

Corollary 4.9. *If $\mathcal{V}$ is a locally presentable monoidal category where $\otimes$ preserves colimits in both entries, the forgetful functors*

$$S : \mathbf{Mon}(\mathcal{V}\text{-}\mathbf{Mat}(X,X)) \to \mathcal{V}\text{-}\mathbf{Mat}(X,X)$$
$$U : \mathbf{Comon}(\mathcal{V}\text{-}\mathbf{Mat}(X,X)) \to \mathcal{V}\text{-}\mathbf{Mat}(X,X)$$

are monadic and comonadic respectively, and all categories are locally presentable.

Notice that $(\mathcal{V}\text{-}\mathbf{Mat}(X,X), \circ, 1_X)$ is a non-symmetric category, therefore the categories of monoids and comonoids cannot inherit a monoidal structure.

4.2. **$\mathcal{V}$-graphs and $\mathcal{V}$-categories.** In this section, we will describe $\mathcal{V}$-graphs and $\mathcal{V}$-categories within the context of $\mathcal{V}$-matrices. This allows us, by dualizing certain arguments, to later construct the category of $\mathcal{V}$-cocategories in a natural way. This is motivated by realizing enriched categories and cocategories as 'many-object' generalizations of monoids and comonoids in a monoidal category: a one-object $\mathcal{V}$-category is an object in $\mathbf{Mon}(\mathcal{V})$. For enriched graphs or categories, usually there are no required assumptions on the monoidal base $\mathcal{V}$ as it is clear from their definition. In this matrices context though, we ask that $\mathcal{V}$ has coproducts preserved by the tensor product

A (small) *$\mathcal{V}$-graph* $\mathcal{G}$ consists of a set of objects $\mathrm{ob}\mathcal{G}$, and for every pair of objects $x, y \in \mathrm{ob}\mathcal{G}$ an object $\mathcal{G}(x,y) \in \mathcal{V}$. If $\mathcal{G}$ and $\mathcal{H}$ are $\mathcal{V}$-graphs, a *$\mathcal{V}$-graph morphism* $F : \mathcal{G} \to \mathcal{H}$ consists of a function $f : \mathrm{ob}\mathcal{G} \to \mathrm{ob}\mathcal{H}$ between their sets of objects, together with arrows $F_{x,y} : \mathcal{G}(x,y) \to \mathcal{H}(fx, fy)$ in $\mathcal{V}$, for each pair of objects x, y in $\mathcal{G}$. These data, with appropriate compositions and identities, form a category $\mathcal{V}$-**Grph**. There is an evident forgetful functor $Q\colon \mathcal{V}\text{-}\mathbf{Grph} \to \mathbf{Set}$ which maps a graph to its set of objects and a graph morphism to its underlying function.

If $\mathcal{V}$-$\mathbb{M}\mathbf{at}$ is the double category of $\mathcal{V}$-matrices, it follows that the category of graphs is precisely that of its endomorphisms as described in Section 3.2, i.e. $\mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1^\bullet = \mathcal{V}\text{-}\mathbf{Grph}$. Indeed, objects are endo-$\mathcal{V}$-matrices $G : X \nrightarrow X$ given by families of objects $\{G(x,x')\}_X$ in $\mathcal{V}$, and morphisms between them are $\alpha_f : G_X \to H_Y$ as in (21), given by arrows $\alpha_{x,x'} : G(x,x') \to H(fx, fx')$ in $\mathcal{V}$ by (33).

Remark 4.10. This viewpoint is very similar to that of [FGK11, Remark 2.5], where it is observed that the category $\mathbf{Grph}_{\mathcal{E}}$ of graphs and graph morphisms internal to a finitely complete $\mathcal{E}$ is identified with the category of endomorphisms and vertical endomorphism maps in the double category $\mathbb{S}\mathbf{pan}_{\mathcal{E}}$, i.e. in our notation $\mathbb{S}\mathbf{pan}^{\bullet}_{\mathcal{E}} = \mathbf{Grph}_{\mathcal{E}}$.

In fact, $\mathcal{V}$-graph morphisms can equivalently be seen as functions $f : X \to Y$ between the sets of objects, equipped with a 2-cell

$$\begin{array}{ccc} & G & \\ X & \Downarrow \phi & X \\ & f^* \circ H \circ f_* & \end{array}$$

in $\mathcal{V}$-$\mathbf{Mat}$, where f_* and f^* are as in (36). This is clear by the following corollary to Proposition 3.15, since $\mathcal{V}$-$\mathbb{M}\mathbf{at}$ is a fibrant double category.

Proposition 4.11. *The category* $\mathcal{V}$-$\mathbf{Grph}$ *is a bifibration over* $\mathbf{Set}$.

The pseudofunctors giving rise to the fibred and opfibred structure are precisely given by (23) in this case,

$$\begin{array}{ccc} \mathscr{M} : \mathbf{Set}^{\mathrm{op}} \longrightarrow \mathbf{Cat}, & & \mathscr{F} : \mathbf{Set} \longrightarrow \mathbf{Cat} \\ X \mapsto \mathcal{V}\text{-}\mathbf{Mat}(X,X) & & X \mapsto \mathcal{V}\text{-}\mathbf{Mat}(X,X) \\ f \downarrow \quad \uparrow f^* \circ - \circ f_* & & f \downarrow \quad \downarrow f_* \circ - \circ f^* \\ Y \mapsto \mathcal{V}\text{-}\mathbf{Mat}(Y,Y) & & Y \mapsto \mathcal{V}\text{-}\mathbf{Mat}(Y,Y) \end{array} \tag{47}$$

and the Grothendieck categories give the following isomorphic characterization of the category of $\mathcal{V}$-graphs.

Lemma 4.12. *The category* $\mathcal{V}$-$\mathbf{Grph}$ *has objects* $(G, X) \in \mathcal{V}$-$\mathbf{Mat}(X,X) \times \mathbf{Set}$ *and arrows* $(\phi, f) : (G, X) \to (H, Y)$ *or equivalently* (ψ, f) *given by*

$$\begin{cases} \phi : G \to f^* H f_* & \text{in } \mathcal{V}\text{-}\mathbf{Mat}(X,X) \\ f : X \to Y & \text{in } \mathbf{Set} \end{cases} \quad \text{or} \quad \begin{cases} \psi : f_* G f^* \to H & \text{in } \mathcal{V}\text{-}\mathbf{Mat}(Y,Y) \\ f : X \to Y & \text{in } \mathbf{Set}. \end{cases}$$

As a sanity check, using (44) we can explicitly compute the composite

$$(f^* H f_*)_{x,x'} = I \otimes (H f_*)_{x,fx'} = I \otimes H_{fx,fx'} \otimes I \tag{48}$$

hence ϕ has components $\phi_{x,x'} : G_{x,x'} \to I \otimes H_{fx,fx'} \otimes I \cong H_{fx,fx'}$. Similarly,

$$(f_* G f^*)_{y,y'} = \sum_{fx=y, fx'=y'} I \otimes G_{x,x'} \otimes I \cong \sum_{fx=y, fx'=y'} G_{x,x'} \tag{49}$$

so $\psi_{y,y'} : \sum_{\substack{fx=y \\ fx'=y'}} I \otimes G_{x,x'} \otimes I \to H_{y,y'}$ which, for fixed $x \in f^{-1}(y)$ and $x' \in f^{-1}(y')$ corresponds uniquely to $\phi_{x,x'}$.

Notice that the above definition is completely in terms of the bicategory $\mathcal{V}$-$\mathbf{Mat}$. Passing from this to the double $\mathcal{V}$-$\mathbb{M}\mathbf{at}^{\bullet}_1$ perspective will be efficient in the proofs that follow.

When $\mathcal{V}$ is braided, $\mathcal{V}$-$\mathbb{M}\mathbf{at}$ is monoidal double by Proposition 4.1 thus its category of endomorphisms is also monoidal by Proposition 3.14: the tensor product is given like in (38) for endo-1-cells and the monoidal unit is the unit $\mathcal{V}$-matrix $I : \{*\} \nrightarrow \{*\}$. Braiding or symmetry is also inherited from $\mathcal{V}$.

Recall from Corollary 4.4 that when $\mathcal{V}$ is furthermore closed with products, the double category of $\mathcal{V}$-matrices is locally monoidal closed. As a corollary to Proposition 3.21, we can deduce that $\mathcal{V}\text{-}\mathbb{M}\mathbf{at}_1^\bullet = \mathcal{V}\text{-}\mathbf{Grph}$ is also monoidal closed, which follows from considering endo-proarrows in the proof of Proposition 4.3.

Proposition 4.13. *Suppose $\mathcal{V}$ is a braided monoidal closed category with products and coproducts. The restriction of (40) on the endomorphism category*

$$H_1^\bullet \colon \mathcal{V}\text{-}\mathbf{Grph}^{\mathrm{op}} \times \mathcal{V}\text{-}\mathbf{Grph} \to \mathcal{V}\text{-}\mathbf{Grph}$$

*mapping $(G_X,\ H_Y)$ to the graph $H_1^\bullet(G,H)_{Y^X}$ given by $H_1^\bullet(G,H)(s,k) := \prod_{x,x'}[G(x,x'), H(sx,kx')]$ for $s,k \in Y^X$ is the internal hom of $\mathcal{V}$-**Grph**.*

Following [KL01, Wol74] or the more general case of bicategory enrichment [BCSW83], we gather some of the main categorical properties of $\mathcal{V}$-**Grph**.

Proposition 4.14.

*(1) $\mathcal{V}$-**Grph** is complete when $\mathcal{V}$ is;*
*(2) $\mathcal{V}$-**Grph** is cocomplete when $\mathcal{V}$ is;*
*(3) [KL01, 4.4] $\mathcal{V}$-**Grph** is locally presentable when $\mathcal{V}$ is.*

Proof. (1) The constructions of limits of enriched graphs are built up from limits in **Set** and $\mathcal{V}$ in a straightforward way.

(2) Suppose F is a diagram of shape $\mathcal{J}$ in $\mathcal{V}$-**Grph**

$$\begin{array}{rccc} F: & \mathcal{J} & \longrightarrow & \mathcal{V}\text{-}\mathbf{Grph} \\ & j & \dashrightarrow & (G_j, X_j) \\ & {\scriptstyle\theta}\downarrow & & \downarrow{\scriptstyle(\psi_\theta, f_\theta)} \\ & k & \dashrightarrow & (G_k, X_k). \end{array} \tag{50}$$

By Lemma 4.12, f_θ is a function between the sets and $(f_\theta)_* G_j (f_\theta)^* \overset{\psi_\theta}{\Rightarrow} G_k$ is a 2-cell in $\mathcal{V}$-**Mat**. The composite $\mathcal{J} \to \mathcal{V}\text{-}\mathbf{Grph} \to \mathbf{Set}$ has a colimiting cocone $(\tau_j : X_j \to X \mid j \in \mathcal{J})$ in **Set**; since $\tau_j = f_\theta \tau_k$ for any $f_\theta : X_j \to X_k$, we have isomorphisms of $\mathcal{V}$-matrices

$$\begin{array}{ccccc} X_j & \xrightarrow{(\tau_j)_*} & X, & \qquad & X \xrightarrow{(\tau_j)^*} X_j \\ & {\scriptstyle(f_\theta)_*}\searrow \quad \overset{\zeta}{\cong} \quad \nearrow{\scriptstyle(\tau_k)_*} & & & {\scriptstyle(\tau_k)^*}\searrow \quad \overset{\xi}{\cong} \quad \nearrow{\scriptstyle(f_\theta)^*} \\ & X_k & & & X_k \end{array}$$

where ζ and ξ are defined as in (45). Now consider the functor

$$\begin{array}{rccc} K: & \mathcal{J} & \longrightarrow & \mathcal{V}\text{-}\mathbf{Mat}(X,X) \\ & j & \dashrightarrow & (\tau_j)_* G_j (\tau_j)^* \cong (\tau_k)_* (f_\theta)_* G_j (f_\theta)^* (\tau_k)^* \\ & {\scriptstyle\theta}\downarrow & & \downarrow{\scriptstyle(\tau_k)_* \psi_\theta (\tau_k)^*} \\ & k & \dashrightarrow & (\tau_k)_* G_k (\tau_k)^* \end{array} \tag{51}$$

which explicitly maps an arrow $\theta : j \to k$ in $\mathcal{J}$ to the composite 2-cell



(52)

The colimit of K is formed pointwise in $[X \times X, \mathcal{V}]$, so there is a colimiting cocone $(\lambda_j : (\tau_j)_* G_j (\tau_j)^* \to G \,|\, j \in \mathcal{J})$. These data allow us to form a new cocone

$$\big((G_j, X_j) \xrightarrow{(\lambda_j, \tau_j)} (G, X) \,|\, j \in \mathcal{J}\big)$$

for the initial diagram F in $\mathcal{V}$-**Grph**, since the pairs (λ_j, τ_j) commute accordingly with the (ψ_θ, f_θ)'s; this cocone can be checked to be colimiting, since τ_j and λ_j are.

(3) Briefly, if $\mathcal{V}$ is a locally λ-presentable category and the set $\mathscr{G}$ of objects constitutes a strong generator of $\mathcal{V}$, it can be shown that the set

$$\{(\bar{G}, 2) \;/\; G \in \mathscr{G} \text{ or } G = 0\}$$

constitutes a strong generator of $\mathcal{V}$-**Grph**, where the graph $(\bar{G}, 2)$ has as set of objects $2 = \{0, 1\}$ and is given by $\{\bar{G}(0,0) = G, \bar{G}(0,1) = \bar{G}(1,0) = \bar{G}(1,1) = 0\}$ in $\mathcal{V}$. Also, this set is λ-presentable in that the functors $\mathcal{V}$-**Grph**$((\bar{G}, 2), -) : \mathcal{V}$-**Grph** $\to$ **Set** preserve λ-filtered colimits. □

Passing on to $\mathcal{V}$-**Cat**, again following the more general [BCSW83], a $\mathcal{V}$-category is defined to be a monad in the bicategory $\mathcal{V}$-**Mat**. Unravelling Definition 2.5, it consists of a set X together with an endoarrow $A : X \nrightarrow X$ (i.e. a $\mathcal{V}$-graph A_X) equipped with two 2-cells, the multiplication and the unit



satisfying the following axioms:





In terms of components, they are given by

$$M_{x,z} \colon \sum_{y \in X} A(x,y) \otimes A(y,z) \to A(x,z) \quad \text{and} \quad \eta_x \colon I \to A(x,x)$$

which are the usual composition law and identity elements. The relations that M and η have to satisfy give the usual associativity and unit axioms.

By Remark 2.9, a monad in a bicategory is the same as a monoid in the appropriate endoarrow hom-category, *i.e.* a $\mathcal{V}$-category A with set of objects X is a monoid in the monoidal category $(\mathcal{V}\text{-}\mathbf{Mat}(X,X),\circ,1_X)$.

A $\mathcal{V}$-functor $F : \mathcal{A} \to \mathcal{B}$ between two $\mathcal{V}$-categories A_X and B_Y is as usual defined as a morphism of graphs $\alpha_f : A_X \to B_Y$ which respects the composition law and the identities. Naturally, one could ask whether this corresponds to the notion of a monad morphism; as will be clear by what follows, this is not the case, see Remark 4.18.

In fact, the category $\mathcal{V}$-**Cat** is fully encompassed as the category of monads $\mathbf{Mnd}(\mathcal{V}\text{-}\mathbb{M}\mathbf{at})$ of Definition 3.9 for the double category of $\mathcal{V}$-matrices. Indeed, objects are monads $A : X \nrightarrow X$ in its horizontal bicategory, and morphisms are arrows between the underlying graphs that respect the structure: writing down what diagrams (22) give in components for $\mathbb{D} = \mathcal{V}\text{-}\mathbb{M}\mathbf{at}$, we end up to the usual axioms

$$\begin{array}{ccc} A(x,y)\otimes A(y,z) & \xrightarrow{M^A_{x,y,z}} & A(x,z) \\ \downarrow{\scriptstyle \alpha_{x,y}\otimes\alpha_{y,z}} & & \downarrow{\scriptstyle \alpha_{x,z}} \\ B(fx,fy)\otimes B(fy,fz) & \xrightarrow[M^B_{fx,fy,fz}]{} & B(fx,fz), \end{array} \qquad \begin{array}{ccc} I & \xrightarrow{\eta_x} & A(x,x) \\ & {\scriptstyle \eta_{fx}}\searrow & \downarrow{\scriptstyle \alpha_{xx}} \\ & & B(fx,fx) \end{array} \tag{53}$$

Due to the fibrant structure of $\mathcal{V}$-$\mathbb{M}\mathbf{at}$, we obtain the following as a corollary to Proposition 3.17.

Proposition 4.15. *The category* $\mathcal{V}$-**Cat** *is a fibration over* **Set**.

The pseudofunctor that gives rise to this fibration is $\mathscr{M}$ from (47), restricted between the categories of monoids of the endo-hom categories. For this to be well-defined, we note the following corollary to Lemma 3.16 for $\mathbb{D} = \mathcal{V}\text{-}\mathbb{M}\mathbf{at}$.

Corollary 4.16. *Let* B_Y *be a* $\mathcal{V}$*-category. For any function* $f : X \to Y$*,*

$$X \xrightarrow{f_*} Y \xrightarrow{B} Y \xrightarrow{f^*} X$$

is a monoid in $\mathcal{V}$-$\mathbf{Mat}(X,X)$, i.e. $(f^*Bf_*)_X$ *is a* $\mathcal{V}$*-category.*

The new composition and unit are given by (28) in this case, i.e. $M' = f^*\left(M\cdot(B\dot{\varepsilon}B)\right)f_*$ and $\eta' = (f^*\eta f_*)\cdot\dot{\eta}$ using pasting operations, where $\dot{\eta},\dot{\varepsilon}$ are the unit and counit of $f_* \dashv f^*$ as in (43).

Once again, the Grothendieck construction that gives rise to the above fibration provide the following isomorphic characterization of the category of enriched categories and functors; compare with Lemma 4.12.

Lemma 4.17. *The objects of* $\mathcal{V}$-**Cat** *are* $(A,X) \in \mathbf{Mon}(\mathcal{V}\text{-}\mathbf{Mat}(X,X)) \times \mathbf{Set}$ *and morphisms are* $(\phi,f) : (A,X) \to (B,Y)$ *where*

$$\begin{cases} \phi : A \to f^*Bf_* & \text{in } \mathbf{Mon}(\mathcal{V}\text{-}\mathbf{Mat}(X,X)) \\ f : X \to Y & \text{in } \mathbf{Set}. \end{cases}$$

Using the mates correspondence for the appropriate 2-cells as in (20) so that ϕ corresponds to $\bar{\phi} : f_* A \Rightarrow B f_*$, we can the $\mathcal{V}$-functors axioms in the form



The mate's components are $\bar{\phi}_{x,y} : I \otimes A(x,x') \to B(fx, fx') \otimes I$ for $x' \in f^{-1}y$, and these diagrams in components agree with (53) up to tensoring with I's and composing with the left and right unit constraints of $\mathcal{V}$.

Remark 4.18. Notice that $(f_*, \bar{\phi})$ constitutes a colax monad functor (Definition 2.7) between the monads A_X and B_Y in the bicategory $\mathcal{V}$-**Mat**. Evidently, however, it is not true that any colax monad functor given by the data



is a $\mathcal{V}$-functor, since not every $S : X \nrightarrow Y$ is of the form f_* for some function $f : X \to Y$. This explains why the category $\mathcal{V}$-**Cat** cannot be characterized as **Mnd**($\mathcal{V}$-**Mat**) for the bicategory, even if they have the same objects. Similar issues were discussed in a bigger depth in [GS13].

This provides a distinct advantage when cosidering categories of monads in double categories rather than in the horizontal bicategory; the notion of a $\mathcal{V}$-functor properly matches the motion of a monad homomorphism in $\mathcal{V}$-$\mathbb{M}$**at**.

As it is well-known, but also deduced from Proposition 3.14, when $\mathcal{V}$ is a braided monoidal category, $\mathcal{V}$-**Cat** is monoidal via the pointwise tensor product (38) like graphs. We now move on to further properties of this category.

Similarly to the free monoid construction on an object in a monoidal $\mathcal{V}$, there is an endofunctor on $\mathcal{V}$-**Grph** inducing the 'free $\mathcal{V}$-category' monad; for the proof below, see also [BCSW83, KL01]. We spell it out in detail in our context, in order to later dualize for $\mathcal{V}$-cocategories.

Proposition 4.19. *Let $\mathcal{V}$ be a monoidal category with coproducts, such that $\otimes$ preserves them on both sides. The forgetful functor $\tilde{S} : \mathcal{V}\text{-}\mathbf{Cat} \to \mathcal{V}\text{-}\mathbf{Grph}$ has a left adjoint $\tilde{L}$, which maps a $\mathcal{V}$-graph $G : X \nrightarrow X$ to the geometric series*

$$\sum_{n\in\mathbb{N}} G^{\otimes n} : X \nrightarrow X.$$

Proof. Recall that by Proposition 4.7, $\mathcal{V}\text{-}\mathbf{Mat}(X,X)$ admits the same class of colimits as $\mathcal{V}$, and also $\otimes = \circ$ preserves them. Hence, the forgetful S from its category of monoids has a left adjoint as in Proposition 2.11

$$\begin{aligned} L :\ \mathcal{V}\text{-}\mathbf{Mat}(X,X) &\longrightarrow \mathbf{Mon}(\mathcal{V}\text{-}\mathbf{Mat}(X,X)) \\ G &\longmapsto \textstyle\sum_{n\in\mathbb{N}} G^n \end{aligned}$$

which by Lemma 4.17 means that this geometric series is a $\mathcal{V}$-category with set of objects X. The claim is that the induced functor

$$\begin{aligned} \tilde{L} :\ \mathcal{V}\text{-}\mathbf{Grph} &\longrightarrow \mathcal{V}\text{-}\mathbf{Cat} \\ (G,X) &\longmapsto (\textstyle\sum_n G^n, X) \end{aligned} \tag{54}$$

is a left adjoint of $\tilde{S}$, i.e. for any $\mathcal{V}$-category B_Y and $\phi_f \colon G_X \to \tilde{S}(B_Y)$ a $\mathcal{V}$-graph morphism, there exists a unique $\mathcal{V}$-functor $H : (\sum_n G^n)_X \to B_Y$ such that

$$\begin{array}{ccc} (G,X) & \xrightarrow{\ \tilde{\eta}\ } & \tilde{S}(\sum\limits_{n\in\mathbb{N}} G^n, X) \\ & {\scriptstyle F}\searrow \quad \swarrow{\scriptstyle \tilde{S}H} & \\ & \tilde{S}(B,Y) & \end{array} \tag{55}$$

commutes, for $\tilde{\eta} : (G,X) \to \tilde{S}\tilde{L}(G,X)$ the identity-on-objects inclusion of the summand G into the series.

By Lemma 4.12, a $\mathcal{V}$-graph functor F is a pair (ϕ, f) with $\phi : G \to f^* B f_*$ an arrow in $\mathcal{V}\text{-}\mathbf{Mat}(X,X)$, and furthermore Corollary 4.16 ensures that $f^* B f_*$ obtains a monoid structure. Since $L(G)$ is the free monoid on $G \in \mathcal{V}\text{-}\mathbf{Mat}(X,X)$, ϕ extends uniquely to a monoid morphism $\chi : LG \to f^* B f_*$ such that

$$\begin{array}{ccc} G & \xrightarrow{\ \eta\ } & \sum\limits_{n\in\mathbb{N}} G^n \\ & {\scriptstyle \phi}\searrow \quad \swarrow{\scriptstyle S\chi} & \\ & f^* B f_* & \end{array}$$

commutes in $\mathcal{V}\text{-}\mathbf{Mat}(X,X)$, where η and S are respectively the unit and forgetful functor of the 'free monoid' adjunction $L \dashv S$. By Lemma 4.17, this 2-cell $\chi : \sum_n G^n \Rightarrow f^* B f_*$ in $\mathcal{V}\text{-}\mathbf{Mat}$ determines a $\mathcal{V}$-functor $H = (\chi, f) : (LG, X) \to (B, Y)$ satisfying the universal property (55). These data suffice to define an adjoint functor $\tilde{L}$ (54), thus the 'free $\mathcal{V}$-category' adjunction $\tilde{L} \dashv \tilde{S} : \mathcal{V}\text{-}\mathbf{Cat} \to \mathcal{V}\text{-}\mathbf{Grph}$ is established. □

Also, as proved in detail in [Wol74] and later generalized in [BCSW83], $\mathcal{V}\text{-}\mathbf{Cat}$ has and the forgetful functor $\tilde{S}$ reflects split coequalizers when $\mathcal{V}$ is cocomplete. By Beck's monadicity theorem, since $\tilde{S}$ also reflects isomorphisms, we have the following well-known result.

Proposition 4.20. *If $\mathcal{V}$ is a cocomplete monoidal category such that $\otimes$ preserves colimits on both variables, $\tilde{S} : \mathcal{V}$-**Cat** $\to \mathcal{V}$-**Grph** is monadic.*

Since $\mathcal{V}$-**Grph** is complete when $\mathcal{V}$ is, we obtain the following corollary.

Corollary 4.21. *The category $\mathcal{V}$-**Cat** is complete when $\mathcal{V}$ is.*

Moreover, by Proposition 4.14 $\mathcal{V}$-**Grph** admits all colimits if $\mathcal{V}$ does; since any category of Eilenberg-Moore algebras has colimits if it has coequalizers of reflexive pairs and its base has colimits by a standard result in [Lin69], the following is also true.

Corollary 4.22. *The category $\mathcal{V}$-**Cat** is cocomplete when $\mathcal{V}$ is.*

Finally, as shown in [KL01] the monad $\tilde{S}\tilde{L}$ is finitary. Thus by [GU71, Satz 10.3] which states that the category of algebras for a finitary monad over a locally presentable category retains that structure, we obtain the following.

Theorem. [KL01, 4.5] *If $\mathcal{V}$ is a monoidal closed category whose underlying ordinary category is locally λ-presentable, then $\mathcal{V}$-**Cat** is also λ-presentable.*

4.3. **$\mathcal{V}$-cocategories.** We now proceed to the dualization of the concept of a $\mathcal{V}$-category in the context of $\mathcal{V}$-matrices, following Definition 2.8. Henceforth $\mathcal{V}$ is again a monoidal category with coproducts, such that the tensor product $\otimes$ preserves them on both entries.

Definition 4.23. A *$\mathcal{V}$-cocategory* C is a comonad in the bicategory $\mathcal{V}$-**Mat**: it consists of a set X with an endoarrow $C : X \nrightarrow X$ (*i.e.* a $\mathcal{V}$-graph C_X) equipped with two 2-cells, the comultiplication and the counit



satisfying the following axioms:



(56)



In terms of components, the *cocomposition* of a $\mathcal{V}$-cocategory C is given by

$$\Delta_{x,z} : C(x,z) \to \sum_{y \in X} C(x,y) \otimes C(y,z)$$

for any two objects $x, y \in X$, and the *coidentity elements* are given by

$$\epsilon_{x,y} : C(x,y) \to 1_X(x,y) \equiv \begin{cases} C(x,x) \xrightarrow{\epsilon_{x,x}} I, & \text{if } x = y \\ C(x,y) \xrightarrow{\epsilon_{x,y}} 0, & \text{if } x \neq y \end{cases}$$

for all objects $x \in X$. The coassociativity and counity axioms are



where α is the associator and λ, ρ are the unitors of $\mathcal{V}$-**Mat**.

As for any comonad in a bicategory, a $\mathcal{V}$-cocategory C with $\mathrm{ob}C = X$ is the same as a comonoid in the monoidal category $(\mathcal{V}\text{-}\mathbf{Mat}(X, X), \circ, 1_X)$. Thus a one-object $\mathcal{V}$-cocategory is the same as a comonoid in $\mathcal{V}$.

A $\mathcal{V}$-cofunctor between two $\mathcal{V}$-cocategories C_X, D_Y should be a $\mathcal{V}$-graph morphism $F_f \colon C_X \to D_Y$ that respects cocomposition and coidentities. As a result, we obtain the following definition.

Definition 4.24. A $\mathcal{V}$-*cofunctor* $F_f : C_X \to D_Y$ between two $\mathcal{V}$-cocategories consists of a function $f : X \to Y$ between their sets of objects and arrows $F_{x,z} : C(x, z) \to D(fx, fz)$ in $\mathcal{V}$ for any $x, z \in \mathrm{ob}C$, satisfying the commutativity of



(57)

Along with compositions and identities of cofunctors that follow those of graphs, we obtain a category $\mathcal{V}$-**Cocat**. As was the case for $\mathcal{V}$-**Cat**, this category is fully encapsulated as the category of comonads in the double category of $\mathcal{V}$-matrices, i.e. $\mathbf{Comon}(\mathcal{V}\text{-}\mathbb{M}\mathbf{at}) = \mathcal{V}\text{-}\mathbf{Cocat}$ as in Definition 3.10. Indeed, an object is just a comonad in the horizontal bicategory, and the comonad homomorphism axioms in components give (57).

Remark 4.25. At [DS97, §9], a $\mathcal{V}$-*opcategory* $\mathcal{A}$ as a category enriched in $\mathcal{V}^{\mathrm{op}}$ is defined. In particular, the cocomposition and counit arrows are simply

$$\Delta \colon \mathcal{A}(x, z) \to \mathcal{A}(y, z) \otimes \mathcal{A}(x, y), \quad \varepsilon \colon \mathcal{A}(x, x) \to I.$$

Also, a $\mathcal{V}$-*opfunctor* $F \colon \mathcal{A} \to \mathcal{B}$ is a $\mathcal{V}^{\mathrm{op}}$-functor, mapping $x \mapsto Fx$ and with arrows $\mathcal{B}(Fx, Fy) \to \mathcal{A}(x, y)$ in $\mathcal{V}$ satisfying respective axioms. The category

$\mathcal{V}$-**opCat** is in a sense broader than $\mathcal{V}$-**Cocat**$^{(\mathrm{op})}$, since for example for $\mathcal{V} = \mathbf{Mod}_R$ any $\mathcal{V}$-cocategory is a special case of a $\mathcal{V}$-category:



i.e. a $\mathbf{Mod}_R$-cocategory is a $\mathbf{Mod}_R$-opcategory for which the cocomposition $\Delta_{x,z,y}$ vanishes for all but finitely many objects z. However, the cocategories point of view seems to be useful in other settings too; in a series of papers related to A_∞-categories, see [Lyu03, Kel06, KM07], the authors study and employ *cocategories* and *cocategory homorphisms* which are precisely our $\mathbf{Mod}_R$-enriched case, in order to express A_∞-functor categories as internal hom objects, building models of a closed structure of the homotopy category of differential graded categories.

Moreover, for the current development it is crucial that a $\mathcal{V}$-cocategory can be expressed as a comonoid in the monoidal endo-hom-categories of $\mathcal{V}$-**Mat**, whereas for a $\mathcal{V}$-opcategory this does not seem to be the case in general.

Due to the fibrant structure of $\mathcal{V}$-$\mathbb{M}$**at**, we again have the following as a corollary to Proposition 3.17.

Proposition 4.26. *The category* $\mathcal{V}$-**Cocat** *is an opfibration over* **Set**.

The pseudofunctor giving rise to this opfibration is $\mathscr{F}$ from (47), now restricted to the categories of comonoids of the endo-hom categories. Again for this to be well-defined, dually to Corollary 4.16 we have the following consequence of $f_* \circ \text{-} \circ f^*$ being colax monoidal.

Lemma 4.27. *Let* C_X *be a* $\mathcal{V}$*-cocategory. If* $f : X \to Y$ *is a function, then*

$$Y \overset{f^*}{\longmapsto} X \overset{C}{\longmapsto} X \overset{f_*}{\longmapsto} Y$$

is a comonoid in $\mathcal{V}$-**Mat**(Y,Y)*, i.e.* $(f_* C f^*)_Y$ *is a* $\mathcal{V}$*-cocategory.*

The new comultiplication and counit come from (29); using pasting operations, we can express them as $\Delta' = f_* \left((C \dot{\eta} C) \cdot \Delta\right) f^*$, $\epsilon' = \dot{\varepsilon} \cdot (f_* \epsilon f^*)$.

The Grothendieck category gives again an isomorphic characterization of the category of $\mathcal{V}$-cocategories and cofunctors, completely in terms of the bicategory $\mathcal{V}$-**Mat**.

Lemma 4.28. *Objects in* $\mathcal{V}$-**Cocat** *are* $(C,X) \in \mathbf{Comon}(\mathcal{V}\text{-}\mathbf{Mat}(X,X)) \times \mathbf{Set}$ *and morphisms are* $(\psi, f) : (C,X) \to (D,Y)$ *where*

$$\begin{cases} \psi : f_* C f^* \to D & \text{in } \mathbf{Comon}(\mathcal{V}\text{-}\mathbf{Mat}(Y,Y)) \\ f : X \to Y & \text{in } \mathbf{Set}. \end{cases}$$

Comparing with the two equivalent formulations for $\mathcal{V}$-graph morphisms of Lemma 4.12, notice that $\mathcal{V}$-functors are expressed as pairs (ϕ, f) and $\mathcal{V}$-cofunctors are expressed as pairs (ψ, f), where the 2-cells $\phi : G \Rightarrow f^* H f_*$ and $\psi : f_* G f^* \Rightarrow H$ are mates under $f_* \dashv f^*$.

Writing explicitly what it means for ψ to be a comonoid morphism, which is clearer in terms of its mate $\hat{\phi} : f_* C \Rightarrow D f_*$, we obtain



(58)

The components of $\hat{\phi}$ are given by $\sum_{x' \in f^{-1} y} I \otimes C(x, x') \to D(fx, fx') \otimes I$ and the above relations componentwise give (57) up to isomorphism. Dually to Remark 4.18, cofunctors correspond to specific types of lax comonad functors in $\mathcal{V}$-**Mat**; viewing them as comonad homomorphisms in $\mathcal{V}$-$\mathbb{M}$**at** is conceptually simpler and less technical, but both expressions will be of use.

When $\mathcal{V}$ is braided monoidal, by Proposition 3.14 $\mathcal{V}$-**Cocat** obtains a monoidal structure, as for $\mathcal{V}$-graphs and categories: for two $\mathcal{V}$-cocategories C_X and D_Y, $(C \otimes D)_{X \times Y}$ is given by $(C \otimes D)\,((x, y), (z, w)) = C(x, z) \otimes D(y, w)$ in $\mathcal{V}$. Writing down in components what the dual of Proposition 3.12 gives for the pseudo double functor $\otimes$ on $\mathcal{V}$-$\mathbb{M}$**at**, we deduce that the cocomposition is

$$\begin{array}{ccc} C(x,z) \otimes D(y,w) & \dashrightarrow & \sum_{(x',y')} C(x,x') \otimes D(y,y') \otimes C(x',z) \otimes D(y',w) \\ & & \uparrow s \cong \\ & & \sum_{(x',y')} C(x,x') \otimes C(x',z) \otimes D(y,y') \otimes D(y',w) \\ & \searrow^{\Delta^C_{x,z} \otimes \Delta^D_{y,w}} & \uparrow \cong \\ & & \sum_{x'} C(x,x') \otimes C(x',z) \otimes \sum_{y'} D(y,y') \otimes D(y',w) \end{array}$$

and the coidentity element is $C(x, x) \otimes D(y, y) \xrightarrow{\epsilon^C_{x,x} \otimes \epsilon^D_{y,y}} I \otimes I \cong I$. The monoidal $\mathcal{V}$-**Cocat** also inherits the braiding or symmetry from $\mathcal{V}$.

Dually to Proposition 4.19, we now construct the cofree $\mathcal{V}$-cocategory functor using the cofree comonoid construction. As discussed in Section 2.2, the existence of the cofree comonoid usually requires more assumptions on $\mathcal{V}$ than the free monoid, and the following is no exception.

Proposition 4.29. *Suppose $\mathcal{V}$ is a locally presentable monoidal category, such that $\otimes$ preserves colimits in both variables. Then, the evident forgetful functor*

$$\tilde{U} : \mathcal{V}\text{-}\mathbf{Cocat} \longrightarrow \mathcal{V}\text{-}\mathbf{Grph}$$

has a right adjoint $\tilde{R}$, which maps a $\mathcal{V}$-graph G_Y to the cofree comonoid (RG, Y) on $G \in \mathcal{V}\text{-}\mathbf{Mat}(Y, Y)$.

Proof. The forgetful $\tilde{U}$ maps any $\mathcal{V}$-cocategory C_X to its underlying $\mathcal{V}$-graph $(UC)_X$, where U is the forgetful from the category of comonoids of the monoidal $(\mathcal{V}\text{-}\mathbf{Mat}(Y, Y), \circ, 1_Y)$. By Corollary 4.9, U has a right adjoint

$$R : \mathcal{V}\text{-}\mathbf{Mat}(Y, Y) \longrightarrow \mathbf{Comon}(\mathcal{V}\text{-}\mathbf{Mat}(Y, Y)),$$

the cofree comonoid functor. By Lemma 4.28, $(RG)_Y$ for any $G : Y \nrightarrow Y$ is a $\mathcal{V}$-cocategory; we claim that

$$\begin{aligned} \tilde{R} : \ \mathcal{V}\text{-}\mathbf{Grph} &\longrightarrow \mathcal{V}\text{-}\mathbf{Cocat} \\ G_Y &\longmapsto (RG)_Y \end{aligned}$$

is a right adjoint of $\tilde{U}$. It suffices to show that for ε the counit of $U \dashv R$, the $\mathcal{V}$-graph arrow $\tilde{\varepsilon} = (\varepsilon, \mathrm{id}_Y) : \tilde{U}\tilde{R}(G_Y) \to G_Y$ is universal. This means that for any $\mathcal{V}$-cocategory C_X and any $\mathcal{V}$-graph morphism F from its underlying $\tilde{U}(C_X)$ to G_Y, there exists a unique $\mathcal{V}$-cofunctor $H : C_X \to (RG)_Y$ such that

$$\begin{array}{ccc} \tilde{U}(RG_Y) & \xrightarrow{\ \tilde{\varepsilon}\ } & G_Y \\ & {\scriptstyle \tilde{U}H} \nwarrow \quad \nearrow {\scriptstyle F} & \\ & \tilde{U}(C_X) & \end{array} \tag{59}$$

commutes. Indeed, suppose F is (ψ, f) with $f : X \to Y$ and $\psi : f_* C f^* \to G$ in $\mathcal{V}\text{-}\mathbf{Mat}(Y, Y)$. Since $f_* C f^*$ is in $\mathbf{Comon}(\mathcal{V}\text{-}\mathbf{Mat}(Y, Y))$ due to Lemma 4.27 and RG is the cofree comonoid on G, ψ extends uniquely to a comonoid arrow $\chi : f_* C f^* \to RG$ such that

$$\begin{array}{ccc} RG & \xrightarrow{\ \varepsilon\ } & G \\ & {\scriptstyle U\chi} \nwarrow \quad \nearrow {\scriptstyle \psi} & \\ & f_* C f^* & \end{array}$$

commutes in $\mathcal{V}\text{-}\mathbf{Mat}(Y, Y)$. By Lemma 4.28, this χ along with the function $f : X \to Y$ determines a $\mathcal{V}$-cofunctor $H : (C, X) \to (RG, Y)$ which makes (59) commute. Therefore $\tilde{R}$ extends to a functor establishing the 'cofree $\mathcal{V}$-cocategory' adjunction $\tilde{U} \dashv \tilde{R} : \mathcal{V}\text{-}\mathbf{Grph} \to \mathcal{V}\text{-}\mathbf{Cocat}$. □

At this point, properties of $\mathcal{V}$-**Cocat** cease to be straightforward dualizations of the $\mathcal{V}$-**Cat** ones. The results that follow more or less employ similar techniques as for the one-object case, that of comonoids in a monoidal category, generalized to this context.

The construction of colimits in $\mathcal{V}$-**Cocat** follows from that in $\mathcal{V}$-**Grph** of Proposition 4.14, with an induced extra structure on the colimiting cocone.

Proposition 4.30. *If $\mathcal{V}$ is a locally presentable monoidal category such that $\otimes$ preserves colimits in both terms, the category $\mathcal{V}$-**Cocat** has all colimits.*

Proof. Consider a diagram in $\mathcal{V}$-**Cocat** given by

$$\begin{array}{rccc} D: & \mathcal{J} & \longrightarrow & \mathcal{V}\text{-}\mathbf{Cocat} \\ & j & \mapsto & (C_j, X_j) \\ & {\scriptstyle\theta}\downarrow & & \downarrow{\scriptstyle(\psi_\theta, f_\theta)} \\ & k & \mapsto & (C_k, X_k) \end{array}$$

for a small category $\mathcal{J}$. By Lemma 4.28, $f_\theta : X_j \to X_k$ is a function and ψ_θ is an arrow $(f_\theta)_* C_j (f_\theta)^* \to C_k$ in $\mathbf{Comon}(\mathcal{V}\text{-}\mathbf{Mat}(X_k, X_k))$. First constructing the colimit of the underlying $\mathcal{V}$-graphs, we obtain a colimiting cocone

$$\big((C_j, X_j) \xrightarrow{(\lambda_j, \tau_j)} (C, X) \,|\, j \in \mathcal{J}\big) \tag{60}$$

in $\mathcal{V}$-**Grph**, where $(\tau_j : X_j \to X \,|\, j \in \mathcal{J})$ is the colimit of the sets of objects of the $\mathcal{V}$-cocategories in **Set**, and $(\lambda_j : (\tau_j)_* C_j (\tau_j)^* \to C \,|\, j \in \mathcal{J})$ is the colimiting cocone of the diagram K as in (51) in the cocomplete $\mathcal{V}$-$\mathbf{Mat}(X, X)$. In fact, $K : \mathcal{J} \to \mathcal{V}\text{-}\mathbf{Mat}(X, X)$ lands inside $\mathbf{Comon}(\mathcal{V}\text{-}\mathbf{Mat}(X, X))$: Lemma 4.27 ensures that $\mathcal{V}$-matrices of the form $f_* C f^*$ for any comonoid C inherit a comonoid structure, and the composite arrows (52) where the middle 2-cell is now the comonoid arrow ψ_θ ensure that $K\theta$ are comonoid morphisms.

By Corollary 4.9, the category of comonoids is comonadic over $\mathcal{V}$-$\mathbf{Mat}(X, X)$ hence the forgetful functor creates all colimits and so $C : X \nrightarrow X$ obtains a unique comonoid structure. Moreover, the legs of the cocone

$$\begin{array}{ccc} X_j & \xrightarrow{C_j} & X_j \\ {\scriptstyle(\tau_j)^*}\uparrow & \Downarrow\lambda_j & \downarrow{\scriptstyle(\tau_j)_*} \\ X & \xrightarrow[C]{} & X \end{array}$$

are comonoid arrows, so together with the functions τ_j they form $\mathcal{V}$-cofunctors. Therefore the colimit (60) lifts in $\mathcal{V}$-**Cocat**. □

We will now apply techniques similar to Proposition 2.13 regarding the expression of the category $\mathbf{Comon}(\mathcal{V})$ as an equifier, to obtain the following.

Proposition 4.31. *Suppose that $\mathcal{V}$ is a locally presentable monoidal category, such that $\otimes$ preserves colimits in both terms. Then, $\mathcal{V}$-**Cocat** is a locally presentable category.*

Proof. Using Lemma 4.12, we can define an endofunctor on $\mathcal{V}$-graphs by

$$\begin{array}{rccc} F: & \mathcal{V}\text{-}\mathbf{Grph} & \longrightarrow & \mathcal{V}\text{-}\mathbf{Grph} \\ & (G, X) & \mapsto & (G \circ G, X) \times (1_X, X) \\ & {\scriptstyle(\psi, f)}\downarrow & & \downarrow{\scriptstyle F(\psi, f)} \\ & (H, Y) & \mapsto & (H \circ H, Y) \times (1_Y, Y) \end{array}$$

with explicit mapping on arrows, for a 2-cell $\psi : f_* G f^* \Rightarrow H$,



(61)

The category of functor coalgebras $\mathbf{Coalg}F$ has as objects $\mathcal{V}$-graphs (C, X) with a morphism $\alpha : C \to C \circ C \times 1_X$, *i.e.* two $\mathcal{V}$-graph arrows

$$\alpha_1 : (C, X) \to (C \circ C, X) \quad \text{and} \quad \alpha_2 : (C, X) \to (1_X, X).$$

A morphism $(C, \alpha) \to (D, \beta)$ is a $\mathcal{V}$-graph morphism $(\psi, f) : (C, X) \to (D, Y)$ which is compatible with α and β, *i.e.* satisfy the equalities





Clearly $\mathbf{Coalg}F$ contains $\mathcal{V}$-$\mathbf{Cocat}$ as a full subcategory: the morphisms satisfy the same axioms (58) for the mate of ψ, and objects are $\mathcal{V}$-graphs equipped with cocomposition and coidentities arrows that don't necessarily satisfy coassociativity and counity.

Since $\mathcal{V}$-$\mathbf{Cocat}$ is cocomplete by Proposition 4.30, we only need to show that it is accessible. It is enough to express it as an equifier of a family of pairs of natural transformations between accessible functors. First of all, F preserves all filtered colimits: take a colimiting cocone for a small filtered category $\mathcal{J}$

$$\big((G_j, X_j) \xrightarrow{(\lambda_j, \tau_j)} (G, X) \,|\, j \in \mathcal{J}\big)$$

in $\mathcal{V}$-$\mathbf{Grph}$ for a diagram like (50) constructed in that proof, *i.e.* $(\tau_j : X_j \to X)$ is colimiting in $\mathbf{Set}$ and $(\lambda_j : (\tau_j)_* C_j (\tau_j)^* \to C)$ is colimiting in $\mathcal{V}$-$\mathbf{Mat}(X, X)$. We require its image under F

$$F(\lambda_j, \tau_j) : (G_j \circ G_j, X_j) \times (1_{X_j}, X_j) \to (G \circ G, X) \times (1_X, X) \tag{62}$$

to be colimiting in $\mathcal{V}$-$\mathbf{Grph}$. For its first part (61), we can deduce that

$$(\tau_j)_* \circ G_j \circ (\tau_j)^* \circ (\tau_j)_* \circ G_j \circ (\tau_j)^* \xrightarrow{\lambda_j * \lambda_j} G \circ G$$

is a colimit in $(\mathcal{V}\text{-}\mathbf{Mat}(X,X), \circ, 1_X)$, as the tensor product (horizontal composite) of two colimiting cocones. Pre-composing this with the unit

$$1 * \dot{\eta} * 1 : (\tau_j)_* \circ G_j \circ 1_{X_j} \circ G_j \circ (\tau_j)^* \to (\tau_j)_* \circ G_j \circ (\tau_j)^* \circ (\tau_j)_* \circ G_j \circ (\tau_j)^*$$

still gives a colimiting cocone: if we take components in $\mathcal{V}$ of the respective 2-cells in $\mathcal{V}$-**Mat**, this comes down to showing that the inclusion

$$\sum_{\substack{z \in X_j \\ \tau_j w = x}}^{\tau_j u = x'} G_j(u,z) \otimes G_j(z,w) \hookrightarrow \sum_{\tau_j a = \tau_j b}^{\substack{\tau_j u = x' \\ \tau_j w = x}} G_j(u,a) \otimes G_j(b,w)$$

for any two fixed $x, x' \in X$, where $u, w, a, b \in X_j$, does not alter the colimit. One way of showing this is by considering the following discrete opfibrations over the filtered shape $\mathcal{J}$:

$$\begin{aligned} \mathcal{L} &= \{(j,a,b) \,|\, j \in \mathcal{J}, a, b \in X_j, \tau_j a = \tau_j b\} \\ \mathcal{M} &= \{(j,z) \,|\, j \in \mathcal{J}, z \in X_j\} \end{aligned}$$

where for example the arrows $(j,a,b) \to (j',a',b')$ in $\mathcal{L}$ are determined by arrows $\theta : j \to j'$ in $\mathcal{J}$ such that $a' = f_\theta(a)$ and $b' = f_\theta(b)$ (the function $f_\theta : X_j \to X_{j'}$ is the image of the diagram (50) in **Set**). We can now define diagrams of shape $\mathcal{L}$ and $\mathcal{M}$ in $\mathcal{V}$

$$\begin{array}{rccc} L: & \mathcal{L} & \longrightarrow & \mathcal{V} \\ & (j,a,b) & \longmapsto & G_j(u,a) \otimes G_j(b,w) \end{array} \qquad \begin{array}{rccc} M: & \mathcal{M} & \longrightarrow & \mathcal{V} \\ & (j,z) & \longmapsto & G_j(u,z) \otimes G_j(z,w) \end{array}$$

and appropriately on morphisms. The colimits for these diagrams in $\mathcal{V}$, taking into account that the fibres are discrete categories, are

$$\begin{aligned} \operatorname{colim} L &\cong \operatorname*{colim}_j \sum_{\tau_j a = \tau_j b} G_j(u,a) \otimes G_j(b,w) \\ \operatorname{colim} M &\cong \operatorname*{colim}_j \sum_{z \in X_j} G_j(u,z) \otimes G_j(z,w). \end{aligned}$$

Finally, notice that there exists a functor $T : \mathcal{M} \to \mathcal{L}$ mapping each (j,z) to (j,z,z) and making the triangle



commute. Due to the construction of filtered colimits in **Set**, it is not hard to show that the slice category $\big((j,z,w) \downarrow T\big)$ is non-empty and connected. Hence T is a final functor and we can restrict the diagram on $\mathcal{L}$ to $\mathcal{M}$ without changing the colimit, as claimed.

For the second part of the diagram (61), it is enough to show that



is a colimiting cocone in $\mathcal{V}$-**Mat**(X,X), for the diagram mapping each j to

$$X \xrightarrow{(\tau_j)^*} X_j \xrightarrow{1_{X_j}} X_j \xrightarrow{(\tau_j)_*} X.$$

This can be established by first verifying that $\dot{\varepsilon}$ is a cocone, and then that it has the required universal property. We have thus shown that the cocone (62) is indeed colimiting, hence F is an accessible functor as required.

Since $\mathcal{V}$-**Grph** is locally presentable and the endofunctor F preserves filtered colimits, **Coalg**F is a locally presentable category by Lemma 2.14 and the forgetful functor $\overline{V} : \mathbf{Coalg}F \to \mathcal{V}\text{-}\mathbf{Grph}$ creates all colimits. We consider the following pairs of transformations between functors from **Coalg**F to $\mathcal{V}$-**Grph**:

$$\phi^1, \psi^1 : \overline{V} \Rightarrow F F \overline{V}, \quad \phi^2, \psi^2 : \overline{V} \Rightarrow (- \circ 1_X)\overline{V}, \quad \phi^3, \psi^3 : \overline{V} \Rightarrow \overline{V}(- \circ 1_X)$$

with natural components



The full subcategory of **Coalg**F spanned by those objects (C,X) which satisfy $\phi^i_{(C,X)} = \psi^i_{(C,X)}$ is precisely the category of $\mathcal{V}$-cocategories by (56), thus

$$\mathbf{Eq}((\phi^i, \psi^i)_{i=1,2,3}) = \mathcal{V}\text{-}\mathbf{Cocat}.$$

Since all categories and functors involved are accessible, $\mathcal{V}$-**Cocat** is accessible. □

The fact that $\mathcal{V}$-**Cocat** is a locally presentable category is very useful for the proof of existence of various adjoints, as seen below.

Proposition 4.32. *Suppose $\mathcal{V}$ is a locally presentable monoidal category such that $\otimes$ preserves colimits in both entries. The forgetful functor $\tilde{U} : \mathcal{V}$-**Cocat** $\to$ $\mathcal{V}$-**Grph** is comonadic.*

Proof. By Proposition 4.29 the forgetful $\tilde{U}$ has a right adjoint, namely the cofree $\mathcal{V}$-cocategory functor $\tilde{R}$. By adjusting (6), consider the commutative



where the top functor is the inclusion in the functor coalgebra category as described above, and the respective forgetful functors discard the structures maps α of the coalgebras. By Lemma 2.14 $\overline{V}$ creates equalizers of split pairs, so it is enough to show that the inclusion ι also creates equalizers of split pairs, since we already have $\tilde{U} \dashv \tilde{R}$. Both $\mathcal{V}$-**Cocat** and $\mathcal{V}$-**Grph** are locally presentable categories so in particular complete, and it is easy to see that ι preserves and reflects, thus creates, all limits. Hence $\tilde{U}$ satisfy the conditions of Precise Monadicity Theorem and the result follows. □

Proposition 4.33. *Suppose that $\mathcal{V}$ is a locally presentable symmetric monoidal closed category. Then the category of $\mathcal{V}$-cocategories is symmetric monoidal closed as well.*

Proof. If $\otimes\colon \mathcal{V}\text{-}\mathbf{Cocat} \times \mathcal{V}\text{-}\mathbf{Cocat} \to \mathcal{V}\text{-}\mathbf{Cocat}$ is the symmetric monoidal structure of $\mathcal{V}$-**Cocat** described earlier, we can form the commutative

$$\begin{array}{ccc} \mathcal{V}\text{-}\mathbf{Cocat} & \xrightarrow{-\otimes D_Y} & \mathcal{V}\text{-}\mathbf{Cocat} \\ {\scriptstyle \tilde{U}}\downarrow & & \downarrow{\scriptstyle \tilde{U}} \\ \mathcal{V}\text{-}\mathbf{Grph} & \xrightarrow[-\otimes \tilde{U}(D_Y)]{} & \mathcal{V}\text{-}\mathbf{Grph} \end{array}$$

where the comonadic $\tilde{U}$ creates all colimits and the bottom arrow preserves them by the monoidal closed structure of $\mathcal{V}$-**Grph**, Proposition 4.13. Therefore $(- \otimes D_Y)$ preserves colimits, and Theorem 2.12 ensures it has a right adjoint since $\mathcal{V}$-**Cocat** is a locally presentable category. If we call it $\textsc{Hom}(D_Y, -)$, we obtain a parametrized adjunction

$$\mathcal{V}\text{-}\mathbf{Cocat} \underset{\textsc{Hom}(D_Y,-)}{\overset{-\otimes D_Y}{\underset{\longleftarrow}{\overset{\longrightarrow}{\bot}}}} \mathcal{V}\text{-}\mathbf{Cocat}$$

which exhibits the uniquely induced $\textsc{Hom}\colon \mathcal{V}\text{-}\mathbf{Cocat}^{\mathrm{op}} \times \mathcal{V}\text{-}\mathbf{Cocat} \to \mathcal{V}\text{-}\mathbf{Cocat}$ as its internal hom. □

4.4. **Enrichment of $\mathcal{V}$-categories in $\mathcal{V}$-cocategories.** Having described the categories of $\mathcal{V}$-graphs, $\mathcal{V}$-categories and $\mathcal{V}$-cocategories in terms of $\mathcal{V}$-matrices, and specified some of their categorical properties relatively to limits and colimits, local presentability and monoidal closed structure, we are now in position to explore an enrichment relation Theorem 4.37 as a many-object generalization of Theorem 2.19 for monoids and comonoids. In particular, viewing $\mathcal{V}$-**Cat** and $\mathcal{V}$-**Cocat** as monads and comonads in the locally symmetric monoidal closed fibrant double category $\mathcal{V}$-$\mathbb{M}$**at**, the desired result will follow from the relevant development in Section 3.3.

Supposed that $\mathcal{V}$ is a braided monoidal closed category with products and coproducts. Recall that the locally monoidal closed structure of the double category of $\mathcal{V}$-matrices established by Corollary 4.4 is given by a lax double functor

$$H = (H_0, H_1)\colon \mathcal{V}\text{-}\mathbb{M}\mathbf{at}^{\mathrm{op}} \times \mathcal{V}\text{-}\mathbb{M}\mathbf{at} \to \mathcal{V}\text{-}\mathbb{M}\mathbf{at}$$

where H_0 is the exponentiation in **Set** and H_1 is defined in (40). This induces a functor **Mon**H (31) as the restriction of $H_1^{\bullet}$ between the category of monads

and comonads, which in this context becomes

$$K\colon \mathcal{V}\text{-}\mathbf{Cocat}^{\mathrm{op}} \times \mathcal{V}\text{-}\mathbf{Cat} \longrightarrow \mathcal{V}\text{-}\mathbf{Cat} \tag{63}$$

mapping a $\mathcal{V}$-cocategory and a $\mathcal{V}$-category (C_X, B_Y) to $K(C,B)_{Y^X}$ given by

$$K(C,B)(s,k) = \prod_{x,x'} [G(x,x'), H(sx, kx')] \text{ for } s,k \in Y^X.$$

Of course this comes from the induced internal hom in $\mathcal{V}$-**Grph**, Proposition 4.13, with the extra structure of a category coming from the more general Proposition 3.12. Explicitly, for each triple $s,k,t \in Y^X$, the composition $M : K(C,B)(s,k) \otimes K(C,B)(k,t) \to K(C,B)(s,t)$ for $\mathcal{K}(C,B)$ is an arrow

$$\prod_{a,a} [C(a,a'), B(sa,ka')] \otimes \prod_{b,b'} [C(b,b'), B(kb,tb')] \to \prod_{c,c'} [C(c,c'), B(sc,tc')].$$

This is defined via its adjunct under the tensor-hom adjunction

$$\begin{array}{l}
\prod\limits_{a,a'} [C_{a,a'}, B_{sa,ka'}] \otimes \prod\limits_{b,b'} [C_{b,b'}, B_{kb,tb'}] \otimes C(c,c') \dashrightarrow B_{sc,tc'} \\
\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad \uparrow M_{sc,tc'} \\
\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad 1\otimes\Delta_{c,c'} \downarrow \\
\prod\limits_{a,a'} [C_{a,a'}, B_{sa,ka'}] \otimes \prod\limits_{b,b'} [C_{b,b'}, B_{kb,tb'}] \otimes \sum\limits_{c''} C_{c,c''} \otimes C_{c'',c'} \qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad \uparrow M_{sc,tc'} \\
\qquad\qquad\qquad\qquad\qquad\qquad\

Proposition 4.35. *Suppose that $\mathcal{V}$ is a locally presentable symmetric monoidal closed category. The functor K^{op} has a parametrized adjoint*

$$T : \mathcal{V}\text{-}\mathbf{Cat}^{\mathrm{op}} \times \mathcal{V}\text{-}\mathbf{Cat} \longrightarrow \mathcal{V}\text{-}\mathbf{Cocat}, \tag{64}$$

given by adjunctions $K(-, B_Y)^{\mathrm{op}} \dashv T(-, B_Y)$ for every $\mathcal{V}$-category B_Y.

Proof. If $H_1^\bullet$ is the internal hom of $\mathcal{V}$-**Grph** as in Proposition 4.13, we can form a square which commutes by definition of K

$$\begin{array}{ccc} \mathcal{V}\text{-}\mathbf{Cocat} & \xrightarrow{K(-,B_Y)^{\mathrm{op}}} & \mathcal{V}\text{-}\mathbf{Cat}^{\mathrm{op}} \\ \downarrow{\scriptstyle \tilde{U}} & & \downarrow{\scriptstyle \tilde{S}} \\ \mathcal{V}\text{-}\mathbf{Grph} & \xrightarrow[H_1^\bullet(-,\tilde{S}B_Y)^{\mathrm{op}}]{} & \mathcal{V}\text{-}\mathbf{Grph}^{\mathrm{op}} \end{array}$$

where the left and right legs create all colimits by Proposition 4.20 and 4.32 and the bottom arrow preserves all colimits by $H_1^\bullet(-, G_Y)^{\mathrm{op}} \dashv H_1^\bullet(-, G_Y)$ in any monoidal closed category. Therefore $K(-, B_Y)^{\mathrm{op}}$ is cocontinuous, and since its domain is locally presentable by Proposition 4.31, Theorem 2.12 provides adjunctions $K(-, B_Y)^{\mathrm{op}} \dashv T(-, B_Y) : \mathcal{V}\text{-}\mathbf{Cat}^{\mathrm{op}} \to \mathcal{V}\text{-}\mathbf{Cocat}$ for all $\mathcal{V}$-categories B_Y; this uniquely induces (64). □

The functor T which generalizes the universal measuring comonoid functor (8) is called the *generalized Sweedler hom*. Morever, the functor K^{op} fixed in the second variable has also a right adjoint, which generalizes the Sweedler product (9).

Lemma 4.36. *There is an adjunction $K(C_X, -)^{\mathrm{op}} \dashv (C_X \triangleright -)^{\mathrm{op}}$, for any $\mathcal{V}$-cocategory C_X.*

Proof. Consider the commutative diagram

$$\begin{array}{ccc} \mathcal{V}\text{-}\mathbf{Cat} & \xrightarrow{K(C_X,-)} & \mathcal{V}\text{-}\mathbf{Cat} \\ \downarrow{\scriptstyle \tilde{S}} & & \downarrow{\scriptstyle \tilde{S}} \\ \mathcal{V}\text{-}\mathbf{Grph} & \xrightarrow[H_1^\bullet(\tilde{U}C_X,-)]{} & \mathcal{V}\text{-}\mathbf{Grph} \end{array}$$

where $\tilde{S}$ is the monadic forgetful functor and the locally presentable category $\mathcal{V}$-**Cat** has all coequalizers. Thus by Dubuc's Adjoint Triangle Theorem [Dub68], the existence of a left adjoint of the bottom arrow by monoidal closedness of $\mathcal{V}$-**Grph** implies the existence of a left adjoint $(C_X \triangleright -)$ of the top arrow. □

The induced functor of two variables $\triangleright : \mathcal{V}\text{-}\mathbf{Cocat} \times \mathcal{V}\text{-}\mathbf{Cat} \to \mathcal{V}\text{-}\mathbf{Cat}$ is called the *generalized Sweedler product*. Now Theorem 3.23 applies to provide the desired enrichment, via the action of the symmetric monoidal closed category $\mathcal{V}$-**Cocat** (Proposition 4.33).

Theorem 4.37. *Suppose $\mathcal{V}$ is a symmetric monoidal closed category which is locally presentable, and T is the generalized Sweedler hom functor.*

(1) $\mathcal{V}$-$\mathbf{Cat}^{\mathrm{op}}$ *is enriched in* $\mathcal{V}$-$\mathbf{Cocat}$*, tensored and cotensored, with hom-objects* $\underline{\mathcal{V}\text{-}\mathbf{Cat}}^{\mathrm{op}}(A_X, B_Y) = T(B_Y, A_X)$.
(2) $\mathcal{V}$-$\mathbf{Cat}$ *is a tensored and cotensored* ($\mathcal{V}$-$\mathbf{Cocat}$)*-enriched category, with*

$$\underline{\mathcal{V}\text{-}\mathbf{Cat}}(A_X, B_Y) = T(A_X, B_Y)$$

cotensor product $K(C,B)_{Yz}$ *and tensor product* $C_Z \triangleright A_X$*, for any* $\mathcal{V}$*-cocategory* C_Z *and any* $\mathcal{V}$*-categories* A_X, B_Y.

The final goal is to combine the above enrichment with the (op)fibrations that these categories form, and characterize them as enriched fibrations, Definition 2.27. First of all, the fibration $P\colon \mathcal{V}\text{-}\mathbf{Cat} \to \mathbf{Set}$ (Proposition 4.15) as well as the opfibration $W\colon \mathcal{V}\text{-}\mathbf{Cocat} \to \mathbf{Set}$ (Proposition 4.26) are both monoidal by Proposition 3.18, due to their expression as categories of monads and comonads in $\mathcal{V}$-$\mathbb{M}\mathbf{at}$; they inherit their symmetry from $\mathcal{V}$. The development of Section 3.3 concluded to Theorem 3.24, which applied in our setting will give the required result.

First of all, we need to show that K^{op} preserves cocartesian liftings, or equivalently K preserves cartesian liftings between the fibrations

$$\begin{array}{ccc} \mathcal{V}\text{-}\mathbf{Cocat}^{\mathrm{op}} \times \mathcal{V}\text{-}\mathbf{Cat} & \xrightarrow{K} & \mathcal{V}\text{-}\mathbf{Cat} \\ \downarrow{\scriptstyle W^{\mathrm{op}}\times P} & & \downarrow{\scriptstyle P} \\ \mathbf{Set}^{\mathrm{op}} \times \mathbf{Set} & \xrightarrow[(-)^{(-)}]{} & \mathbf{Set}. \end{array} \tag{65}$$

Using the canonical (co)cartesian liftings (11) for any Grothendieck fibration, i.e. $\mathrm{Cocart}(f, C) = (1_{f_* C f^*}, f)\colon B \to f_* C f^*$ for a cocategory C_X and $f\colon X \to Z$, $\mathrm{Cart}(g, B) = (1_{g^* B g_*}, g)\colon g^* B g_* \to B$ for a category B_Y and $g\colon W \to Y$, we want to deduce that K maps them to the chosen cartesian lifting

$$\begin{array}{ccc} K(f_* C f^*, g^* B g_*) & \xrightarrow{K(\mathrm{Cart}(f,C),\mathrm{Cart}(g,B))} & K(C,B) \\ \cong\downarrow & \nearrow_{\mathrm{Cart}(g^f, K(C,B))} & \vdots \\ (g^f)^* K(C,B)(g^f)_* & & \vdots \\ \downarrow & & \vdots \\ W^Z & \xrightarrow[g^f]{} & Y^X \end{array} \qquad \begin{array}{l} \\ \\ \text{in } \mathcal{V}\text{-}\mathbf{Cat} \\ \\ \text{in } \mathbf{Set} \end{array}$$

Using (48) and (49), we can initially compute

$$\begin{aligned} K(f_* C f^*, g^* B g_*)_{s,k} &= \prod_{z,z' \in Z} [(f_* C f^*)_{z,z'}, (g^* B g_*)_{sz,kz'}] \\ &\cong \prod_{z,z' \in Z} [\sum_{fx=z}^{fx'=z'} C_{x,x'}, B_{gsz,gkz'}], \\ \big((g^f)^* K(C,B)(g^f)_*\big)_{s,k} = K(C,B)_{gsf,gsk} &= \prod_{x,x' \in X} [C_{x,x'}, B_{gsfx,gsfx'}] \end{aligned}$$

which are isomorphic since the internal hom maps sums to products in the first variable, and the triangle commutes by also applying K to the maps.

The only thing left to show is that the opposite of (65) has a general lax parametrized opfibred adjoint, i.e. the opfibred 1-cell for any $\mathcal{V}$-category B_Y

$$\begin{array}{ccc} \mathcal{V}\text{-}\mathbf{Cocat} & \xrightarrow{K(-,B_Y)^{\mathrm{op}}} & \mathcal{V}\text{-}\mathbf{Cat}^{\mathrm{op}} \\ {\scriptstyle W}\downarrow & & \downarrow{\scriptstyle P^{\mathrm{op}}} \\ \mathbf{Set} & \xrightarrow[Y^{(-)^{\mathrm{op}}}]{} & \mathbf{Set}^{\mathrm{op}} \end{array}$$

has a lax opfibred adjoint; this will be deduced from Theorem 2.23. Indeed, there is an adjunction between the base categories

$$\mathbf{Set} \underset{Y^{(-)}}{\overset{Y^{(-)^{\mathrm{op}}}}{\underset{\longleftarrow}{\overset{\longrightarrow}{\bot}}}} \mathbf{Set}^{\mathrm{op}}$$

since $\mathbf{Set}$ is cartesian monoidal closed, with counit ε. Moreover, we need to show is that the composite (12) between the fibres

$$\mathcal{V}\text{-}\mathbf{Cocat}_{Y^Z} \xrightarrow{K_{Y^Z}(-,B_Y)^{\mathrm{op}}} \mathcal{V}\text{-}\mathbf{Cat}^{\mathrm{op}}_{Y^{Y^Z}} \xrightarrow{(\varepsilon_Z)_!} \mathcal{V}\text{-}\mathbf{Cat}^{\mathrm{op}}_{Z}$$

has a right adjoint. We can rewrite it as

$$\begin{array}{ccc} \mathbf{Comon}(\mathcal{V}\text{-}\mathbf{Mat}(Y^Z,Y^Z)) & \xrightarrow{\mathbf{Mon}(\mathrm{Hom}(-,B_Y))^{\mathrm{op}}} & \mathbf{Mon}(\mathcal{V}\text{-}\mathbf{Mat}(Y^{Y^Z},Y^{Y^Z}))^{\mathrm{op}} \\ & \searrow & \downarrow{\scriptstyle (\varepsilon)^* \circ - \circ (\varepsilon)_*} \\ & & \mathbf{Mon}(\mathcal{V}\text{-}\mathbf{Mat}(Z,Z))^{\mathrm{op}} \end{array}$$

where the top functor is (46) between the categories of monoids (as a restriction of $\mathbf{Mon}H$ on globular 2-cells) and the side functor is the reindexing functor for the fibration P, Proposition 4.15. By Corollary 4.9, the domain $\mathbf{Comon}(\mathcal{V}\text{-}\mathbf{Mat}(Y^Z,Y^Z))$ is a locally presentable category for any set Y^Z. Moreover, the following commutative

$$\begin{array}{ccc} \mathbf{Comon}(\mathcal{V}\text{-}\mathbf{Mat}(Y^Z,Y^Z)) & \xrightarrow{\mathbf{Mon}(\mathrm{Hom}(-,B_Y)^{\mathrm{op}}} & \mathbf{Mon}(\mathcal{V}\text{-}\mathbf{Mat}(Y^{Y^Z},Y^{Y^X}))^{\mathrm{op}} \\ {\scriptstyle U}\downarrow & & \downarrow{\scriptstyle S^{\mathrm{op}}} \\ \mathcal{V}\text{-}\mathbf{Mat}(Y^Z,Y^Z) & \xrightarrow[\mathrm{Hom}(-,B_Y)^{\mathrm{op}}]{} & \mathcal{V}\text{-}\mathbf{Mat}(Y^{Y^Z},Y^{Y^Z})^{\mathrm{op}} \end{array}$$

for a fixed $\mathcal{V}$-category B_Y shows that the top arrow is cocontinuous: U and S^{op} are comonadic by the same Corollary 4.9 and the bottom arrow is the cocontinuous internal hom of $\mathcal{V}$-$\mathbf{Grph}$ (Proposition 4.13) restricted between the cocomplete fibres. Finally, composing with the companion and conjoint of ε on either side always preserves colimits, since tensoring does (see Proposition 4.7). Therefore, Theorem 2.12 establishes an adjunction

$$\mathcal{V}\text{-}\mathbf{Cocat}_{Y^Z} \underset{T_0(-,B_Y)}{\overset{(\varepsilon_Z)_! \circ K(-,B_Y)^{\mathrm{op}}}{\underset{\longleftarrow}{\overset{\longrightarrow}{\bot}}}} \mathcal{V}\text{-}\mathbf{Cat}^{\mathrm{op}}_{Z}$$

between the fibre categories, enough by Theorem 2.23 to induce a right adjoint of $K(-, B_Y)^{\mathrm{op}}$ between the total categories such that

$$\begin{array}{ccc} \mathcal{V}\text{-}\mathbf{Cocat} & \underset{T(-,B_Y)}{\overset{K(-,B_Y)^{\mathrm{op}}}{\rightleftarrows}} & \mathcal{V}\text{-}\mathbf{Cat}^{\mathrm{op}} \\ {\scriptstyle W}\downarrow & & \downarrow{\scriptstyle P^{\mathrm{op}}} \\ \mathbf{Set} & \underset{Y^{(-)}}{\overset{Y^{(-)^{\mathrm{op}}}}{\rightleftarrows}} & \mathbf{Set}^{\mathrm{op}} \end{array}$$

is a general lax opfibred adjunction. The assumptions of Theorem 3.24 are now satisfied and the result follows.

Theorem 4.38. *Suppose $\mathcal{V}$ is a symmetric monoidal closed category, which is locally presentable. The opfibration $\mathcal{V}$-**Cat**$^{\mathrm{op}} \to$ **Set**$^{\mathrm{op}}$ as well as the fibration $\mathcal{V}$-**Cat** $\to$ **Set** are enriched in the symmetric monoidal opfibration $\mathcal{V}$-**Cocat** $\to$ **Set**.*

Notice that the total parametrized adjoint $T\colon \mathcal{V}\text{-}\mathbf{Cat}^{\mathrm{op}} \times \mathcal{V}\text{-}\mathbf{Cat} \to \mathcal{V}\text{-}\mathbf{Cocat}$ of K obtained as above is isomorphic to (64), but the fibred approach provided with the extra information that the underlying set of objects of some $T(A_X, B_Y)$ is precisely Y^X in a straightforward way.

References

[AJ13] Matthieu Anel and André Joyal. Sweedler theory of (co)algebras and the bar-cobar constructions. arXiv:1309.6952 [math.CT], 2013.

[AR94] Jiří Adámek and Jiří Rosický. *Locally presentable and accessible categories*, volume 189 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1994.

[BCSW83] Renato Betti, Aurelio Carboni, Ross Street, and Robert Walters. Variation through enrichment. *J. Pure Appl. Algebra*, 29(2):109–127, 1983.

[BCV16] E. Batista, S. Caenepeel, and J. Vercruysse. Hopf categories. *Algebr. Represent. Theory*, 19(5):11731216, 2016.

[Bén67] Jean Bénabou. Introduction to bicategories. In *Reports of the Midwest Category Seminar*, pages 1–77. Springer, Berlin, 1967.

[Bén73] J. Bénabou. *Les distributeurs*. Rapport (Université catholique de Louvain (1970-).33 Séminaire de mathématique pure)). Institut de mathématique pure et appliquée, Université catholique de Louvain, 1973.

[BL16] Gabriella Bhm and Stephen Lack. Hopf comonads on naturally frobenius map-monoidales. *Journal of Pure and Applied Algebra*, 220(6):2177 – 2213, 2016.

[Bor94a] Francis Borceux. *Handbook of categorical algebra. 1*, volume 50 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1994. Basic category theory.

[Bor94b] Francis Borceux. *Handbook of categorical algebra. 2*, volume 51 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1994. Categories and structures.

[BS76] Ronald Brown and Christopher B. Spencer. Double groupoids and crossed modules. *Cahiers Topologie Géom. Différentielle*, 17(4):343–362, 1976.

[B16] Gabriella Bhm. Hopf polyads, hopf categories and hopf group monoids viewed as hopf monads. arXiv:1611.05157 [math.CT], 2016.

[Car95] Sean Carmody. *Cobordism categories*. PhD thesis, University of Cambridge, 1995.

[DPP10] Robert Dawson, Robert Paré, and Dorette Pronk. The span construction. *Theory Appl. Categ.*, 24:No. 13, 302–377, 2010.

[DS97] Brian Day and Ross Street. Monoidal bicategories and Hopf algebroids. *Adv. Math.*, 129(1):99–157, 1997.

[Dub68] Eduardo Dubuc. Adjoint triangles. In *Reports of the Midwest Category Seminar, II*, pages 69–91. Springer, Berlin, 1968.

[Ehr63] Charles Ehresmann. Catégories structurées. *Ann. Sci. École Norm. Sup. (3)*, 80:349–426, 1963.

[FGK11] Thomas M. Fiore, Nicola Gambino, and Joachim Kock. Monads in double categories. *J. Pure Appl. Algebra*, 215(6):1174–1197, 2011.

[FGK12] Thomas M. Fiore, Nicola Gambino, and Joachim Kock. Double adjunctions and free monads. *Cah. Topol. Géom. Différ. Catég.*, 53(4):242–306, 2012.

[GP97] R. Gordon and A. J. Power. Enrichment through variation. *J. Pure Appl. Algebra*, 120(2):167–185, 1997.

[GP99] Marco Grandis and Robert Paré. Limits in double categories. *Cahiers Topologie Géom. Différentielle Catég.*, 40(3):162–220, 1999.

[GP04] Marco Grandis and Robert Paré. Adjoint for double categories. Addenda to: "Limits in double categories" [Cah. Topol. Géom. Différ. Catég. **40** (1999), no. 3, 162–220; mr1716779]. *Cah. Topol. Géom. Différ. Catég.*, 45(3):193–240, 2004.

[GS13] Richard Garner and Michael Shulman. Enriched categories as a free cocompletion. arXiv:1301.3191 [math.CT], 2013.

[GU71] P. Gabriel and F. Ulmer. *Lokal Präsentierbare Kategorien*, volume 221 of *Lecture Notes in Mathematics*. Springer-Verlag, 1971.

[Her93] Claudio Alberto Hermida. *Fibrations, logical predicates and indeterminates*. PhD thesis, University of Edinburgh, 1993.

[Her94] Claudio Hermida. On fibred adjunctions and completeness for fibred categories. In *Recent trends in data type specification (Caldes de Malavella, 1992)*, volume 785 of *Lecture Notes in Comput. Sci.*, pages 235–251. Springer, Berlin, 1994.

[HLFV15] Martin Hyland, Ignacio Lopez Franco, and Christina Vasilakopoulou. Hopf measuring comonoids and enrichment. arXiv:1509.07632 [math.CT], 2015.

[HLFV17] Martin Hyland, Ignacio Lopez Franco, and Christina Vasilakopoulou. Measuring comodules and enrichment. arXiv:1703.10137 [math.CT], 2017.

[Jac99] Bart Jacobs. *Categorical logic and type theory*, volume 141 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, 1999.

[JK02] G. Janelidze and G. M. Kelly. A note on actions of a monoidal category. *Theory Appl. Categ.*, 9:61–91, 2001/02. CT2000 Conference (Como).

[JS93] André Joyal and Ross Street. Braided tensor categories. *Adv. Math.*, 102(1):20–78, 1993.

[Kel05] G. M. Kelly. Basic concepts of enriched category theory. *Repr. Theory Appl. Categ.*, (10):vi+137, 2005. Reprint of the 1982 original [Cambridge Univ. Press, Cambridge; MR0651714].

[Kel06] Bernhard Keller. *A*-infinity algebras, modules and functor categories. In *Trends in representation theory of algebras and related topics*, volume 406 of *Contemp. Math.*, pages 67–93. Amer. Math. Soc., Providence, RI, 2006.

[KL01] G. M. Kelly and Stephen Lack. V-Cat is locally presentable or locally bounded if V is so. *Theory Appl. Categ.*, 8:555–575, 2001.

[KM07] Bernhard Keller and Oleksandr Manzyuk. Equalizers in the category of cocomplete cocategories. *J. Homotopy Relat. Struct.*, 2(1):85–97, 2007.

[KS74] G. Kelly and Ross Street. Review of the elements of 2-categories. In Gregory Kelly, editor, *Category Seminar*, volume 420 of *Lecture Notes in Mathematics*, pages 75–103. Springer Berlin / Heidelberg, 1974. 10.1007/BFb0063101.

[Lac10] Stephen Lack. A 2-categories companion. In *Towards higher categories*, volume 152 of *IMA Vol. Math. Appl.*, pages 105–191. Springer, New York, 2010.

[Lin69] F. E. J. Linton. Coequalizers in categories of algebras. In *Sem. on Triples and Categorical Homology Theory (ETH, Zürich, 1966/67)*, pages 75–90. Springer, Berlin, 1969.

[LV07] Jean-Louis Loday and Bruno Vallette. *Algebraic Operads*. Grundlehren der mathematischen Wissenschaften. Springer, 2007.

[Lyu03] Volodymyr Lyubashenko. Category of A_∞-categories. *Homology Homotopy Appl.*, 5(1):1–48, 2003.

[ML98] Saunders Mac Lane. *Categories for the working mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1998.

[MP89] Michael Makkai and Robert Paré. *Accessible categories: the foundations of categorical model theory*, volume 104 of *Contemporary Mathematics*. American Mathematical Society, Providence, RI, 1989.

[Por08a] Hans-E. Porst. Fundamental constructions for coalgebras, corings, and comodules. *Appl. Categ. Structures*, 16(1-2):223–238, 2008.

[Por08b] Hans-E. Porst. On categories of monoids, comonoids, and bimonoids. *Quaest. Math.*, 31(2):127–139, 2008.

[Shu08] Michael Shulman. Framed bicategories and monoidal fibrations. *Theory Appl. Categ.*, 20:No. 18, 650–738, 2008.

[Shu10] Michael Shulman. Constructing symmetric monoidal bicategories. arXiv:1004.0993 [math.CT], 2010.

[Str72] Ross Street. The formal theory of monads. *J. Pure Appl. Algebra*, 2(2):149–168, 1972.

[Str80] Ross Street. Fibrations in bicategories. *Cahiers Topologie Géom. Différentielle*, 21(2):111–160, 1980.

[Str96] Ross Street. Categorical structures. In *Handbook of algebra, Vol. 1*, volume 1 of *Handb. Algebr.*, pages 529–577. Elsevier/North-Holland, Amsterdam, 1996.

[Str07] Ross Street. *Quantum groups*, volume 19 of *Australian Mathematical Society Lecture Series*. Cambridge University Press, Cambridge, 2007. A path to current algebra.

[Swe69] Moss E. Sweedler. *Hopf algebras*. Mathematics Lecture Note Series. W. A. Benjamin, Inc., New York, 1969.

[Vas14] Christina Vasilakopoulou. *Generalization of Algebraic Operations via Enrichment*. PhD thesis, University of Cambridge, 2014. arxiv:1411.3038 [math.CT].

[Vas17] Christina Vasilakopoulou. Enriched fibrations. in preparation, 2017.

[Ver11] Dominique Verity. Enriched categories, internal categories and change of base. *Theory Appl. Categ.*, pages 1–266, 2011.

[Wol74] Harvey Wolff. V-cat and V-graph. *J. Pure Appl. Algebra*, 4:123–135, 1974.

[Woo82] R. J. Wood. Abstract pro arrows i. 23(3):279–290, 1982.

Département de Mathématiques, Faculté des sciences, Université Libre de Bruxelles, Boulevard du Triomphe, B-1050 Bruxelles, Belgium

E-mail address: christina.vasilakopoulou@ulb.ac.be